

Association of Complement and Coagulation Pathway Proteins With Treatment Response in First-Episode Psychosis: A Longitudinal Analysis of the OPTiMiSE Clinical Trial

Subash Raj Susai^{*,1,a}, Melanie Föcking^{1,a}, David Mongan¹, Meike Heurich², Fiona Coutts³, Alice Egerton^{3,6}, Tony Whetton⁴, Inge Winter-van Rossum⁵, Richard D. Unwin⁶, Thomas A. Pollak^{7,8}, Mark Weiser^{8,9}, Marion Leboyer¹⁰, Dan Rujescu^{11,12}, Jonah F Byrne^{1,6}, George W Gifford³, Paola Dazzan³, Nikolaos Koutsouleris^{7,12}, René S. Kahn¹³, David R. Cotter^{1,b}, and Philip McGuire^{3,b}

¹Department of Psychiatry, RCSI University of Medicine and Health Sciences, Dublin, Ireland; ²School of Pharmacy and Pharmaceutical Sciences, Cardiff University, Cardiff, UK; ³Department of Psychosis Studies, Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London, UK; ⁴School of Veterinary Medicine, School of Biosciences and Medicine, University of Surrey, Guildford, GU2 7XH, UK; ⁵Department of Psychiatry, University Medical Center Utrecht Brain Center, Utrecht University, Utrecht, The Netherlands; ⁶Stoller Biomarker Discovery Centre and Division of Cancer Sciences, School of Medicine, Faculty of Biology Medicine and Health, The University of Manchester, Manchester, M13 9NY, UK; ⁷Institute of Psychiatry, Psychology and Neuroscience, Department of Psychosis Studies, King's College London, London, UK; ⁸Stanley Medical Research Institute, Kensington, MD, USA; ⁹Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel; ¹⁰Université Paris Est Creteil (UPEC), AP-HP, Hôpitaux Universitaires « H. Mondor », DMU IMPACT, FHU ADA-T, INSERMU955, IMRB, Translational Neuropsychiatry laboratory, F-94010 Creteil, France; ¹¹University Clinic and Outpatient Clinic for Psychiatry, Psychotherapy and Psychosomatics, Martin Luther University of Halle-Wittenberg, Halle, Germany; ¹²Department of Psychiatry and Psychotherapy, Ludwig-Maximilian-University, Munich, Germany; ¹³Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA

^aJoint first authors.

^bJoint senior authors.

*To whom correspondence should be addressed; 1. Subash Raj Susai, Department of Psychiatry, RCSI University of Medicine and Health Sciences, Dublin, Ireland; tel: +353 1 809 3862, e-mail: subashrajsusai@rcsi.ie, susaisubash@rcsi.com

Background and Hypothesis: Treatment response to specific antipsychotic medications is difficult to predict on clinical grounds alone. The current study hypothesizes that the baseline complement pathway activity predicts the treatment response and investigates the relationship between baseline plasma biomarkers with treatment response to antipsychotic medications. **Study Design:** Baseline plasma samples were collected from first episode of psychosis patients ($n = 243$) from a multi-center clinical trial. The participants were treated with amisulpride for 4 weeks. Levels of complement and coagulation proteins at baseline were measured using both data-dependent and data-independent mass spectrometry approaches. The primary outcome was remission status at 4 weeks and the secondary outcomes included change in psychotic and functional symptoms over the period of treatment. In addition, immunoassays were performed at baseline for complement C1R, as well as for activation markers C4a and sC5b-9. **Study Results:** The plasma level of complement variant C4A

was significantly associated with remission at 4 weeks. Moreover, higher levels of several complement and coagulation pathway proteins were associated with a reduction in psychotic symptoms and an improvement in functioning. Immunoassays showed an association of baseline levels of C1R and C4a as well as complement activation marker sC5b-9 levels with treatment response. **Conclusion:** The results demonstrated that the response to antipsychotic treatment might be related to pre-treatment levels of plasma complement and coagulation pathway proteins. This is consistent with independent evidence associating immune dysfunction with the pathophysiology of psychosis. Moreover, these results inform the development of novel therapeutic approaches that target the complement system for psychosis.

Key words: Schizophrenia/psychotic disorder/immune markers/biomarkers/complement system proteins/antipsychotic agents

Introduction

Around a third of patients with psychosis do not respond to initial treatment with antipsychotic medication.¹⁻³ Because the response to treatment is difficult to predict based on the clinical presentation, the effectiveness of antipsychotic medication in an individual patient has to be determined empirically, through the careful evaluation of a course of treatment. However, recent research suggests that it may be possible to predict the therapeutic response by assessing biomarkers, measured using neuroimaging⁴⁻⁷ or in peripheral blood.⁸⁻¹⁰

The role of immune system dysfunction in psychosis is increasingly acknowledged.¹¹⁻¹⁷ More specifically, there is some evidence that response to antipsychotic medication may be related to the plasma levels of inflammatory markers^{8,10,18} and auto-antibodies¹⁹ prior to treatment. Immune dysfunction may also be manifested in changes in the concentration of proteins of the complement system.^{20,21} Complement activity in both the peripheral and the central nervous system is crucial to the overall immune response. Complement proteins are mostly present in the circulation as inactive zymogens, which are converted into active enzymes and proteolytic fragments acting as opsonins or chemotactic molecules. This is triggered through the activation of 1 of the 3 complement pathways: the classical, lectin, and alternative pathways (figure 1). These pathways converge in a terminal common pathway that results in formation of the membrane attack complex (MAC or C5b-9). Complement C1R is one of the initiating serine proteases of the C1 complex, crucial for propagation of the classical complement pathway leading to C4 cleavage into proteolytic fragments C4b (opsonin) and C4a. Sequential proteolytic cleavage of C4 and C2 allows the formation of the C3 convertase (C4b2a), which cleaves C3, and if left unregulated, C5.²⁰ This generates quantifiable activation




Fig. 1. Group difference in pre-treatment complement C4A levels between patients who were in remission, or not in remission after 4 weeks of treatment with amisulpride.

products, including C4a, the anaphylatoxins C3a and C5a, and terminal pathway activation marker sC5b-9.²² C4 exists as 2 isotypes, C4A and C4B. A Genome-wide association study reported that the C4 alleles of the major histocompatibility complex were linked to an increased risk of psychosis, and that this risk was proportionate to the expression of C4A.²³⁻²⁸ We recently found that altered plasma levels of complement and coagulation pathway proteins, which are closely connected to each other, were associated with the onset of psychosis in people at clinical high-risk.²⁹⁻³¹

The aim of the present study was to investigate the relationship between levels of complement and coagulation proteins and the response to treatment with antipsychotic medication. By conducting the study in patients with first-episode psychosis who were either medication-naïve or had minimal previous exposure to antipsychotic medications, the potentially confounding effects of previous treatment were minimized. Moreover, because all the patients were treated with the same antipsychotic medication (amisulpride) in a clinical trial, differences in response were unlikely to be due to variation in the type or dose of treatment. We previously performed a pilot study based on a subset of participants in the OPTiMiSE trial (optimization of treatment and management of schizophrenia in Europe) using data-dependent acquisition (DDA) discovery-based proteomic analysis of plasma samples.³² Comparing responders to non-responders, there was preliminary evidence for dysregulation of several complement and coagulation proteins in relation to responders.

In the current study, we tested the hypothesis that the levels of complement and coagulation proteins prior to treatment would be associated with the subsequent response to treatment, as indexed by remission status and the change in symptom severity. To test our hypothesis, we undertook 2 complementary mass spectrometry approaches. First, we undertook a discovery-based approach called “data-dependent acquisition” (DDA), to extend our earlier pilot study³² to the whole OPTiMiSE population and to confirm the findings using the same mass spectrometric methods as previously. Second, to investigate associations between proteins and symptomatic and treatment response, we used a data-independent acquisition (DIA) mass spectrometry approach to allow us to extract data on complement and coagulation pathway proteins in a more targeted manner.

Materials and Methods

Participants

The patients were participants in the OPTiMiSE study, a multi-center trial of amisulpride conducted at 20 institutions and 15 countries (Clinicaltrials.gov identifier NCT01248195). The trial protocol has been described in detail elsewhere.³³ Five-hundred patients with the

first episode of schizophrenia, schizophreniform disorder, or schizoaffective disorder as defined by the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) were verified with the Mini International Neuropsychiatric Interview Plus.³⁴ Participants were aged between 18 and 40 years, had a maximum interval of 2 years between the onset of psychosis and study entry, and had a history of antipsychotic medication of no longer than 2 weeks in the previous year or 6 weeks lifetime.³³ Ethical approval for the proteomic analyses was granted by the Research Ethics Committee of the Royal College of Surgeons Ireland [REC No: 1699].

Outcomes

We examined 3 outcomes. The primary outcome was symptomatic remission as defined by Andreasen et al.³⁵ Participants were assessed for remission after treatment for 4 weeks. In secondary analyses, we examined 2 further outcomes: a change in Positive and Negative Symptom scale (PANSS)³⁶ score from baseline to 4 weeks and a change in Personal and Social Performance Scale (PSP)³⁷ score from baseline to 4 weeks.

Mass Spectrometry Based DDA and DIA Proteomics

Two different mass spectrometry (MS) based proteomic methods were used to relatively quantify the levels of plasma proteins.

First, data-dependent acquisition (DDA) was performed to measure the relative abundance of complement and coagulation proteins in the immunodepleted plasma. Second, the proteins were quantified using targeted extraction from data-independent acquisition (DIA), which was based on sequential window acquisition of all theoretical mass spectra (SWATH) analysis in which all peptide precursor ions in the scanning interval are scanned at ultra-high speed and subjected to secondary fragmentation.^{38,39}

The protocols for these 2 methods are described in detail in the [supplementary methods](#).

Enzyme-Linked Immunoassay (ELISA) Quantifying Complement Protein C1R, and Activation Markers C4a and sC5b-9

We used standardized enzyme-linked immunosorbent assays (ELISA) to validate key mass spectrometry findings and measure the impact on complement activation by quantifying plasma concentrations of C1R (ABCAM Ltd, USA), and downstream activation markers C4a (Abbexa Ltd, UK) and sC5b-9 (Pathway Diagnostics Ltd, UK). C4a is proportional to C4 activation and sC5b-9 measures complement activation of the common terminal pathway (see [supplementary methods](#)).

Statistical Analyses

Statistical analyses were conducted using IBM SPSS statistics version 26 and STATA IC version 16. The set of complement and coagulation proteins that were analyzed were selected based on the UNIPROT functional annotation of pathway proteins. The LFQ values of plasma proteins that were expressed in at least 70% of the samples were considered for the statistical analysis. The mass spectrometry proteomic data were log₂ transformed and standardised to z-scores for further analysis.

Associations Between Protein Levels and Remission Status We first tested whether the levels of complement and coagulation proteins at baseline were associated with remission status (remission/non-remission) after 4 weeks of treatment. For this analysis, baseline standardized plasma protein levels were measured using both DDA and DIA proteomic approaches. The proteomic data were used as the independent variable. For both DDA and DIA data, associations with remission status at 4 weeks were evaluated for each protein using logistic regression models adjusted for age, sex, BMI, smoking status (current smoker/past smoker/no smoker), and study site. *P*-values were corrected for multiple comparisons using the Benjamini-Hochberg method with a false discovery rate (FDR) of 5%.⁴⁰ A receiver operator curve (ROC) analysis was performed for the significant protein C4A to assess its diagnostic value to predict remission status in FEP participants.

Association Between Plasma Protein Levels and Changes in Symptoms and Functioning To evaluate associations between individual protein levels at baseline and changes in psychotic symptoms and functioning, we used the absolute protein levels from the DIA-SWATH proteomics as exposures. Linear regression analyses were performed using changes in PANSS total score and PSP score between baseline and 4 weeks as outcomes and each protein as the exposure. Models were adjusted for age, sex and BMI, smoking status, study site, and baseline PANSS/PSP score. *P*-values were corrected for multiple comparisons using the Benjamini-Hochberg method with a false discovery rate (FDR) of 5%.⁴⁰

Enzyme-Linked Immunoassay (ELISA) Quantifying Complement Protein C1R, and Activation Markers C4a and sC5b-9

Finally, we examined associations between log-transformed standardized concentrations of the complement component C1R, and activation fragments C4a and sC5b-9, as measured by ELISA, and the outcomes measures (see [supplementary methods](#)).

Results

Complement and Coagulation Proteins From DDA and DIA Proteomic Analysis

Of the 534 participants initially recruited, a total of 294 participants had provided baseline samples and had also completed Phase I of the clinical trial. This subgroup was considered for the present study and the proteomic analysis was performed in baseline plasma samples from 294 participants. After the exclusion of participants who already met remission criteria at baseline, 243 participants with complete baseline and follow-up clinical data comprised the analytical sample. The baseline characteristics of the sample are given in [table 1](#).

Data-dependent acquisition (DDA) mass spectrometry identified a total of 455 plasma proteins in all samples. Of these, 111 proteins passed the quality control (identified in more than 70% of plasma samples) and the missing values were imputed using minimum LFQ values of corresponding protein. Out of 111 proteins, 34 proteins belonged to the complement and coagulation cascade activity (proteins of interest) and these proteins were considered for statistical analysis ([supplementary table 1](#)). Using DIA proteomics (SWATH), a total of 648 proteins were qualified (based on the coefficient of variation filtering). Since the DIA quantification is reproducible, the samples with poor co-efficient of variation (CVs) were rerun and the raw data were normalized in MSSStats data analysis pipeline (see [supplementary methods](#)). Out of 648 proteins, 48 proteins belonged to the complement and coagulation cascade (proteins of interest). The baseline plasma levels of these proteins were further used for statistical analysis ([supplementary table 2](#)).

Associations With Remission Status After 4 Weeks of Treatment

Data-Dependent Acquisition (DDA) In the discovery proteomic approach (DDA data), complement components C1R, C2, and coagulation factor XII (F12) were associated with remission status at 4 weeks with odds ratios (95% CI) of 1.42 (1.08, 1.88), 1.47 (1.04–1.88), and 1.40 (1.04, 1.93), respectively. However, these associations were not significant after adjusting for multiple corrections ([supplementary table 1](#)).

Data-Independent Acquisition (DIA) In logistic regression models of the DIA proteomic data, baseline plasma levels of complement protein C4A associated with the primary outcome (remission status) after 4 weeks of treatment, with an odds ratio of 1.60 [CI = 1.20, 2.14] (FDR adjusted *p*-value .046) ([supplementary table 2](#)). The group difference between remitters and non-remitters for C4A is shown in [figure 1](#). In the ROC analysis, baseline levels of C4A indicated a prediction of remission status with an area under the curve (AUC) of 63% (*p* = .001) ([supplementary figure 1](#)). Other complement and coagulation proteins at baseline did not show any association with remission status ([supplementary table 2](#)).

Associations With Changes in Psychotic Symptoms

In linear regression models using DIA data, the levels of 16 proteins at baseline were significantly associated (after FDR correction) with a reduction in total PANSS score over the first 4 weeks of treatment ([table 2](#)). These comprised 11 complement proteins (complement C1R, C1S, C2, C5, C6, C7, C8A, C8B, CFI, FCN3, and MASP1),

Table 1. Baseline Characteristics of a Sub-group of Participants Who Showed Remission (Remitters) and Did Not Show Remission (Non-Remitters) to Amisulpride From the OPTiMiSE Trial

Baseline Characteristics		Non-Remitters	Remitters
		(<i>n</i> = 78)	(<i>n</i> = 165)
Age in years, mean ± SD		24 ± 5	27 ± 7
Sex	Female, <i>n</i> (%)	58 (74.4%)	110 (66.7%)
	Male, <i>n</i> (%)	20 (25.6%)	55 (33.3%)
BMI in kg/m ² , mean ± SD		23.76 ± 4.62	23.16 ± 3.95
Smoking status	Never smoked, <i>n</i> (%)	27 (34.6%)	67 (40.6%)
	Current smoker, <i>n</i> (%)	45 (57.7%)	75 (45.5%)
	Past smoker, <i>n</i> (%)	6 (7.7%)	22 (13.3%)
PANSS, mean ± SD	Total	85.65 ± 16.45	78.98 ± 17.66
	Positive	21.33 ± 5.32	20.45 ± 5.28
	Negative	23.03 ± 7.27	19.3 ± 6.87
	General	41.29 ± 8.71	39.22 ± 9.35
Personal and Social Performance Scale, mean ± SD		45 ± 15	47 ± 14
Calgary Depression Scale, mean ± SD		13.84 ± 4.56	13.58 ± 4.68

Note: SD, Standard deviation; BMI, body mass index; PANSS, positive and negative symptom score.

Table 2. Results of Linear Regression Model Showing Association of Complement and Coagulation Proteins With Change in Psychopathology Scores After 4 Weeks of Treatment

Outcome Protein Names	Change in Total PANSS					Change in PSP Scores				
	Coef.	P Value	[95% Conf. Interval]	FDR p value		Coef.	P value	[95% Conf. Interval]	FDR p value	
Mannan-binding lectin serine protease	1.36	.18	-0.63	3.35	.26	-0.35	.78	-2.32	1.62	.84
Ficolin-3	-4.11	.00*	-6.20	-2.02	.00*	2.82	.01*	0.73	4.91	.06
Coagulation factor XIII A chain	1.56	.12	-0.41	3.54	.20	0.04	.96	-1.88	1.97	.96
Prothrombin	-2.21	.03*	-4.20	-0.22	.07	1.68	.09	-0.29	3.65	.22
Complement CIR subcomponent	-4.39	.00*	-6.35	-2.44	.00*	3.31	.00*	1.37	5.25	.02*
Coagulation factor IX	-3.73	.00*	-5.69	-1.77	.00*	2.36	.02*	0.41	4.32	.09
Coagulation factor X	-3.73	.00*	-5.69	-1.76	.00*	2.98	.00*	1.09	4.88	.02*
Complement factor D	-1.50	.14	-3.48	0.47	.21	-0.14	.89	-2.06	1.79	.93
Coagulation factor XII	-0.68	.51	-2.71	1.34	.63	1.00	.32	-0.97	2.98	.48
Complement factor B	-1.90	.07	-3.94	0.15	.15	2.06	.04*	0.06	4.06	.12
Complement C3	-1.02	.33	-3.07	1.04	.43	0.45	.66	-1.54	2.43	.84
Complement C5	-2.61	.01*	-4.60	-0.62	.03*	1.79	.07	-0.18	3.76	.18
Fibrinogen alpha chain	-0.45	.66	-2.48	1.58	.73	0.30	.76	-1.64	2.23	.84
Fibrinogen beta chain	-0.38	.72	-2.40	1.65	.75	0.12	.91	-1.83	2.06	.93
Fibrinogen gamma chain	-0.48	.64	-2.52	1.56	.73	0.46	.64	-1.49	2.41	.84
Complement C1q subcomponent subunit A	-0.55	.59	-2.56	1.46	.70	-0.34	.74	-2.32	1.64	.84
Complement C1q subcomponent subunit B	0.41	.69	-1.60	2.42	.74	-1.17	.23	-3.08	0.74	.44
Complement C1q subcomponent subunit C	-1.70	.09	-3.69	0.29	.17	0.98	.32	-0.97	2.92	.48
Complement component C9	-1.76	.08	-3.75	0.24	.17	1.07	.28	-0.89	3.03	.47
Coagulation factor XI	-1.70	.10	-3.71	0.30	.17	0.68	.49	-1.27	2.62	.67
Vitronectin	-1.29	.20	-3.29	0.71	.28	2.16	.03*	0.20	4.11	.11
Complement factor I	-3.62	.00*	-5.62	-1.62	.00*	3.45	.00*	1.52	5.39	.02*
Coagulation factor XIII B chain	1.05	.31	-0.96	3.06	.41	-0.37	.71	-2.33	1.59	.84
Heparin cofactor 2	-3.46	.00*	-5.45	-1.47	.00*	3.70	.00*	1.77	5.63	.00*
Complement C2	-2.98	.00*	-4.94	-1.02	.01*	1.60	.10	-0.33	3.52	.23
Complement component C8 alpha chain	-4.05	.00*	-6.01	-2.10	.00*	2.44	.02*	0.48	4.40	.09
Complement component C8 beta chain	-3.02	.00*	-4.98	-1.06	.01*	1.33	.18	-0.62	3.28	.37
Complement component C8 gamma chain	0.09	.93	-1.90	2.07	0.93	1.19	.22	-0.70	3.08	.43
Complement factor H	-1.83	.09	-3.92	0.27	0.17	1.19	.25	-0.83	3.20	.45
Complement C1s subcomponent	-3.29	.00*	-5.28	-1.31	0.00*	2.11	.04*	0.15	4.08	.12
Complement C4-A	-2.34	.02*	-4.31	-0.37	0.05	1.97	.04*	0.07	3.87	.12
Complement C4-B	-1.62	.12	-3.66	0.41	0.20	-0.73	.46	-2.67	1.20	.65
Complement component C7	-2.48	.01*	-4.44	-0.52	0.04*	1.05	.29	-0.89	2.99	.47
Clusterin	-2.35	.02*	-4.35	-0.35	0.06	2.82	.01*	0.87	4.77	.05*
Mannose-binding protein C	1.37	.18	-0.62	3.37	0.26	-1.00	.31	-2.94	0.93	.48
Coagulation factor V	1.69	.10	-0.31	3.68	0.17	-1.96	.04*	-3.83	-0.09	.12
Complement component C6	-3.98	.00*	-5.89	-2.06	0.00*	2.18	.03*	0.25	4.12	.10
Fibulin-1	-2.79	.01*	-4.76	-0.82	0.02*	1.85	.06	-0.06	3.76	.15
Complement factor H-related protein 2	-0.13	.90	-2.13	1.87	0.92	0.30	.77	-1.66	2.26	.84
Mannan-binding lectin serine protease 1	-2.93	.00*	-4.92	-0.93	0.01*	2.33	.02*	0.40	4.25	.09
Complement factor H-related protein 1	-0.81	.42	-2.79	1.16	0.53	0.32	.74	-1.60	2.24	.84

Downloaded from https://academic.oup.com/schizophreniabulletin/advance-article/doi/10.1093/schbul/sbac201/7077182 by Martin-Luther-Universität Halle-Wittenberg user on 12 May 2023

Table 2. Continued

Outcome Protein Names	Change in Total PANSS					Change in PSP Scores				
	Coef.	<i>P</i> Value	[95% Conf. In- terval]		FDR <i>p</i> value	Coef.	<i>P</i> value	[95% Conf. In- terval]		FDR <i>p</i> value
Multimerin-1	-0.58	.57	-2.59	1.42	0.69	0.25	.80	-1.68	2.19	.85
Ficolin-2	0.43	.67	-1.54	2.40	0.73	0.70	.46	-1.19	2.59	.65
Complement factor H-related protein 5	1.98	.05	-0.02	3.97	0.12	-2.31	.02*	-4.23	-0.39	.09
Complement C1R subcomponent-like protein	-1.46	.15	-3.43	0.52	0.23	1.06	.27	-0.83	2.95	.47
<i>Protein Z-dependent pro- tease inhibitor</i>	-3.29	.00*	-5.23	-1.35	0.00*	2.23	.02*	0.33	4.14	.09

Note: The results of significant proteins from the complement and coagulation pathway were marked bold and italic. FDR-False discovery rate, *significant, PANSS, Positive and Negative Symptom Scale; PSP, Personal and Social Performance scale.

and 5 coagulation pathway proteins (coagulation factor F9, F10, FBLN1, Serine protease inhibitor SERPINA10, and SERPIND1).

Associations With Change in Functional Status

In linear regression models using DIA data, increased expression of baseline levels of 5 proteins were associated with increased PSP score from baseline to 4 weeks of treatment (table 2): C1R (β 3.3, 95% CI 1.37–5.35, $p = .015$); CFI (β 3.5, 95% CI 1.52–5.39, $p = .015$); Clusterin (β 2.8, 95% CI 0.87–4.77, $p = .05$); Coagulation factor F10 (β 2.9, 95% CI 1.09–4.88); and SERPIND1 (β 3.7, 95% CI 1.77–5.63, $p < .001$).

Enzyme-Linked Immunoassay (ELISA) Results

In a logistic regression model, the baseline C4a activation marker concentration was significantly associated with remission status [Odds ratio (CI) = 1.48 (1.08–2.04); p -value = .02] (table 3). This is in alignment with the proteomics C4A findings, which also detected an increased expression of C4A associated with remission status. In the proteomic analysis, the high levels of protein C1R at baseline showed a significant association with a reduction in PANSS score. Using ELISA quantification, we observed a similar trend of association between baseline C1R concentration and a reduction in PANSS score (coef = -0.27, 95% CI = -0.54, 0.001, $p = .05$) (supplementary figure 2). The sC5b-9 concentration was inversely associated with an increase in PANSS score (coef = 0.28, 95% CI = 0.03, 0.53, $p = .03$) (table 3).

Discussion

The present study examined the relationship between complement and coagulation proteins and the response to treatment in psychosis. We tested the hypothesis that the levels of these proteins at baseline would be associated with the response to subsequent treatment with

antipsychotic medication. Our results from both DDA and DIA approaches provided results pointing to a similar trend indicating an association between increased baseline levels of complement and coagulation proteins with treatment response to antipsychotics. The DDA proteomic results of complement and coagulation proteins indicated that higher levels of C1R, C2, and F12 were associated with remission (supplementary table 1), but these findings did not survive adjustment for multiple comparisons. Qualitatively similar findings were evident with a DIA proteomic analysis (supplementary table 2), which identified a strong association between higher levels of C4A with remission following amisulpride treatment in FEP. The association between C4A variant levels and remission was further evaluated by quantifying C4 activation fragment C4a, which was similarly associated with remission, indicating an association between elevated C4 activation with treatment response in FEP. In a ROC analysis, C4A indicated a diagnostic property with an AUC value of 63% for predicting remission status in FEP. Collectively, these results suggest that elevated baseline levels of several complement and coagulation proteins are associated with a positive early clinical response to amisulpride.

The DIA proteomic investigation focused on the complement and coagulation pathways and showed an association for 19 complement and coagulation proteins with the change in PANSS-rated symptom score over 4 weeks of treatment. Out of these, 16 proteins (C1R, C1S, C2, C5, C6, C7, C8A, C8B, CFI, FCN3, MASP1, F9, F10, FBLN1, SERPINA10, and SERPIND1) demonstrated a significant association after FDR correction. Similarly, increased expression of 17 complement and coagulation proteins was associated with improvement in PSP-rated functioning over the same period. Of these proteins, the association of 5 proteins (C1R, CFI, CLU, F10, and SERPIND1) remained significant after adjusting for multiple comparisons. Among the proteins that were associated with treatment response, the classical complement protein C1R demonstrated a strong association




Fig. 2. Schematic representation of complement cascade proteins dysregulated at baseline in participants with a poor symptomatic response to treatment Created with BioRender.com.

with both, a decrease in psychotic symptoms and an improvement in functioning. This was confirmed in the C1R ELISA (supplementary figure 2).

Our findings showed that increased levels of both the activators and some inhibitors of the complement cascade are associated with symptomatic improvement in FEP patients (Figure 2). For instance, C1R contributes to the activation of the classical complement pathway, while complement factor I (CFI) is an integral complement inhibitor acting across all pathways.^{41,42} Clusterin is a crucial terminal pathway inhibitor of the terminal product of the complement pathway called C5b-9. The fluid phase sC5b-9 which was quantified in our study is a biomarker of overall complement activity.⁴² In contrast to other complement pathway proteins, the terminal activator complex sC5b-9 levels were significantly lower at baseline among subjects who responded well to amisulpride. Although we observe comparatively lower sC5b-9 levels, the absence of healthy controls limits us from understanding whether the changes in complement activation markers in relation to treatment response is a return to, or a further deviation from, normal physiological limits. Further mechanistic studies are required to elucidate the precise sequence of these changes.

Other than the biological significance, the results carry a potential clinical relevance in the prediction of remission after antipsychotic medication. The ROC analysis of baseline levels of complement protein C4A indicated a weak prediction of remission status (63% AUC) at follow-up after amisulpride treatment. However, the association of several complement cascade proteins with symptomatic response suggests that future studies with a combined complement score calculated from multiple complement proteins could have more predictive value than single biomarkers for the prediction of treatment response in FEP. Several studies have investigated the role of peripheral complement pathway proteins in the pathophysiology of psychosis^{43,44} including the involvement of C4A, which has been associated with schizophrenia risk in genome-wide association studies.^{23,24} To the best of our knowledge, the present study is the first to examine the relationship of several complement cascade proteins with symptomatic response to antipsychotic medication. A recent study by Mondelli et al. in patients with psychosis found that a higher baseline C4 level was associated with the number of relapses of psychosis in FEP at one-year follow-up.⁹ Other studies have investigated associations

Table 3. Results of Logistic Regression Models for Immune-Assays of C5b-9, C1R, and C4a (Immuno-Assay Data) With the Treatment Response

Proteins at Baseline	Outcome	Odds Ratio	<i>p</i> -value	[95% Conf. Interval]	
<i>C4a</i>	Remission status	<i>1.48</i>	<i>.016*</i>	<i>1.08</i>	<i>2.04</i>
C1R		1.11	.496	0.83	1.48
C5b-9		1.02	.896	0.76	1.36

Proteins at Baseline	Outcome	Coefficient	<i>p</i> -value	[95% Conf. Interval]	
C4a	Change in PANSS total score	-0.051	.69	-0.30	0.20
C1R		-0.268	.05	-0.54	0.001
<i>C5b-9</i>		<i>0.279</i>	<i>.03*</i>	<i>0.026</i>	<i>0.532</i>

Proteins at Baseline	Outcome	Coefficient	<i>p</i> -value	[95% conf. Interval]	
C4a	Change in PSP total score	-0.089	.47	-0.327	0.150
<i>C1R</i>		0.053	.69	-0.211	0.317
<i>C5b-9</i>		0.060	.64	-0.188	0.308

Note: The results of significant proteins from the complement and coagulation pathway were marked bold and italic. For C5b-9 and C1R the participants were grouped based on the change PANSS score from baseline to 4 weeks follow-up. The odds ratio was obtained based on the association of baseline proteins predicting the group with higher reduction in PANSS score. For C4a the association of baseline protein levels were analyzed against the remission status based on the Andreasen criteria. C, complement; PANSS, Positive and Negative Symptom Scale.

between other peripheral immune markers and the response to treatment in psychosis. In the OPTiMiSE cohort, Martinuzzi et al. found an association between non-remission after 4 weeks of treatment and lower levels of the cytokine interleukin (IL)-15, but higher levels of the C-X-C motif chemokine 12 (CXCL12).⁴⁵ They also reported that in participants who were medication-naïve, plasma neurotrophin levels were associated with poor functional status after follow-up at 1-year, suggesting the existence of different pathological mechanisms in relation to antipsychotic medication.⁴⁶ Similarly, lower cortisol-awakening response and higher cytokine levels (IL-6 and TNF- α) have been linked with a poor response to antipsychotic treatment after 12 weeks,⁸ whereas serum levels of NMDA receptor auto-antibodies do not appear to predict treatment response.⁴⁷

Our study has several strengths. Investigating first-episode patients with minimal previous exposure to treatment with antipsychotic medication reduced the risk that the response to the trial medication was confounded by the effects of previous treatment or chronic illness. Moreover, all patients were treated with the same antipsychotic medication, following the same dose regimen, so differences in response are unlikely to have been related to variations in the type or dose of medication. We used 2 independent, state-of-the-art mass spectrometry-based approaches (DDA-based discovery and DIA-SWATH semi-targeted analysis) that provided complementary results. In the current study, we restricted the MS-based proteomic analysis to samples that passed stringent quality criteria filters, excluding

16% of the original plasma samples. Several findings were validated using ELISA-based methods.

The study has several limitations. The 2 proteomic methods used slightly different methods of plasma protein depletion, and the influence of such differences in methodology on relative abundance of valuable proteins is not addressed.⁴⁸⁻⁵⁰ In addition, the large-scale discovery proteomic analysis did not find an association of complement protein C4A variant levels and treatment response, despite there being a strong association found in the DIA analysis. C4A, which was identified in the discovery of MS data in more than 40% of the samples, did not pass the stringent quality criteria applied, and therefore the label-free quantification intensities (normalized measures) of C4A were not considered for the DDA data analysis. Finally, the present study examined group-level differences in biomarkers and is not able to address whether proteomic measures can be used to make predictions about treatment response at the individual level. We plan to investigate this in subsequent work.

Conclusion

We report evidence for an association of increased levels of several complement and coagulation proteins in plasma with early clinical response to amisulpride in first-episode psychosis. These results are consistent with evidence that immune dysfunction contributes to the pathophysiology of psychosis and may inform the development of novel therapeutic approaches for psychosis that target the complement system.

Supplementary Material

Supplementary material is available at <https://academic.oup.com/schizophreniabulletin/>.

Acknowledgments

We are extremely grateful to the participants of the OPTiMiSE study. We acknowledge Ms. Magdalena Hryniewiecka for her technical support in preparation of plasma samples for DDA proteomics. We acknowledge the contribution of Prof. Gerard Cagney and Dr. Eugene Dillon at the UCD Conway Mass Spectrometry Core Facility for help with MS experiments. DM, MC, and DRC report a patent pending (UK Patent Application No. 1919155.0, “Biomarkers to predict psychosis”). RDU is an inventor named in patent applications that describe the use of complement inhibitors for therapeutic purposes and the use of circulating complement protein measurement for patient stratification, and is co-founders of and shareholder in Complement Therapeutics, a company which focuses on the development of complement targeted therapeutics for chronic diseases focused on Age-related Macular Degeneration. Other authors declare no conflict of interest in relation to work described in this study.

Funding

The OPTiMiSE clinical trial was funded by European Commission within the 7th Program (HEALTH-F2-2010-242114). The research work was funded also by Irish Health Research Board grant to David Cotter (HRB-ILP-2019-005), Wellcome Trust IMPETUS Innovations Flagship grant (project 220438Z/20/Z). D.M. is a Fellow on the Irish Clinical Academic Training (ICAT) Programme which is supported by the Wellcome Trust and the Health Research Board (Grant Number 203930/B/16/Z), the Health Service Executive National Doctors Training and Planning and the Health and Social Care, Research and Development Division, Northern Ireland. S.R.S is supported by the Health Research Board (HRB) under grant number HRB/HRA/PHR/2015-1293. D.R.C and J.F.B are supported in part by Science Foundation Ireland (SFI) under Grant Number 16/RC/3948 and co-funded under the European Regional Development Fund and by FutureNeuro industry partners.

References

- Meltzer HY. Treatment-resistant schizophrenia—the role of clozapine. *Curr Med Res Opin.* 1997;14(1):1–20.
- Dempster K, Li A, Sabesan P, Norman R, Palaniyappan L. Treatment resistance: a time-based approach for early identification in first episode psychosis. *J Pers Med.* 2021;11(8):711.
- Samara MT, Nikolakopoulou A, Salanti G, Leucht S. How many patients with schizophrenia do not respond to antipsychotic drugs in the short term? An analysis based on individual patient data from randomized controlled trials. *Schizophr Bull.* 2019;45(3):639–646.
- Sarpal DK, Argyelan M, Robinson DG, et al. Baseline striatal functional connectivity as a predictor of response to antipsychotic drug treatment. *Am J Psychiatry.* 2015;173(1):69–77.
- Han S, Becker B, Duan X, et al. Distinct striatum pathways connected to salience network predict symptoms improvement and resilient functioning in schizophrenia following risperidone monotherapy. *Schizophr Res.* 2020;215:89–96.
- Ebdrup BH, Skimminge A, Rasmussen H, et al. Progressive striatal and hippocampal volume loss in initially antipsychotic-naïve, first-episode schizophrenia patients treated with quetiapine: relationship to dose and symptoms. *Int J Neuropsychopharmacol.* 2011;14(1):69–82.
- Oh S, Kim M, Kim T, Lee TY, Kwon JS. Resting-state functional connectivity of the striatum predicts improvement in negative symptoms and general functioning in patients with first-episode psychosis: a 1-year naturalistic follow-up study. *Aust N Z J Psychiatry.* 2020;54(5):509–518.
- Mondelli V, Ciufolini S, Belvederi Murri M, et al. Cortisol and inflammatory biomarkers predict poor treatment response in first episode psychosis. *Schizophr Bull.* 2015;41(5):1162–1170.
- Mondelli V, Di Forti M, Morgan BP, Murray RM, Pariante CM, Dazzan P. Baseline high levels of complement component 4 predict worse clinical outcome at 1-year follow-up in first-episode psychosis. *Brain Behav Immun.* 2020;88:913–915.
- Kose M, Pariante CM, Dazzan P, Mondelli V. The role of peripheral inflammation in clinical outcome and brain imaging abnormalities in psychosis: a systematic review. *Front Psychiatry.* 2021;12:612471.
- Muller N, Weidinger E, Leitner B, Schwarz MJ. The role of inflammation in schizophrenia. *Front Neurosci.* 2015;9:372.
- Müller N. Inflammation in schizophrenia: pathogenetic aspects and therapeutic considerations. *Schizophr Bull.* 2018;44(5):973–982.
- Khandaker GM, Cousins L, Deakin J, Lennox BR, Yolken R, Jones PB. Inflammation and immunity in schizophrenia: implications for pathophysiology and treatment. *Lancet Psychiatry.* 2015;2(3):258–270.
- Khandaker GM, Dantzer R. Is there a role for immune-to-brain communication in schizophrenia? *Psychopharmacology (Berl).* 2016;233(9):1559–1573.
- Buckley PF. Neuroinflammation and Schizophrenia. *Curr Psychiatry Rep.* 2019;21(8):72.
- Pedraz-Petrozzi B, Elyamany O, Rummel C, Mulert C. Effects of inflammation on the kynurenine pathway in schizophrenia—a systematic review. *J Neuroinflammation.* 2020;17(1):56.
- Fond G, Lançon C, Korchia T, Auquier P, Boyer L. The role of inflammation in the treatment of schizophrenia. *Front Psychiatry.* 2020;11:160.
- Mongan D, Sabherwal S, Susai SR, Föcking M, Cannon M, Cotter DR. Peripheral complement proteins in schizophrenia: a systematic review and meta-analysis of serological studies. *Schizophr Res.* 2020;222:58–72.
- Pollak TA, Rogers JP, Nagele RG, et al. Antibodies in the diagnosis, prognosis, and prediction of psychotic disorders. *Schizophr Bull.* 2019;45(1):233–246.

20. Merle NS, Church SE, Fremeaux-Bacchi V, Roumenina LT. Complement system part I - molecular mechanisms of activation and regulation. *Front Immunol.* 2015;6:262.
21. Merle NS, Noe R, Halbwachs-Mecarelli L, Fremeaux-Bacchi V, Roumenina LT. Complement system part II: role in immunity. *Front Immunol.* 2015;6:257.
22. Ekdahl KN, Persson B, Mohlin C, Sandholm K, Skattum L, Nilsson B. Interpretation of serological complement biomarkers in disease. *Front Immunol.* 2018;9:2237.
23. Sekar A, Bialas AR, de Rivera H, et al; Schizophrenia Working Group of the Psychiatric Genomics Consortium. Schizophrenia risk from complex variation of complement component 4. *Nature.* 2016;530(7589):177–183.
24. Sekar A, Bialas A, De Rivera H, et al. Complex structural variation in the MHC locus influences schizophrenia risk by shaping expression of complement component 4. *Eur Neuropsychopharmacol.* 2017;27:S142.
25. Hamshere ML, Walters JT, Smith R, et al; Schizophrenia Psychiatric Genome-wide Association Study Consortium. Genome-wide significant associations in schizophrenia to ITIH3/4, CACNA1C and SDCCAG8, and extensive replication of associations reported by the Schizophrenia PGC. *Mol Psychiatry.* 2013;18(6):708–712.
26. Purcell SM, Moran JL, Fromer M, et al. A polygenic burden of rare disruptive mutations in schizophrenia. *Nature.* 2014;506(7487):185–190.
27. Shi J, Levinson DF, Duan J, et al. Common variants on chromosome 6p22.1 are associated with schizophrenia. *Nature.* 2009;460(7256):753–757.
28. Irish Schizophrenia Genomics Consortium and the Wellcome Trust Case Control Consortium 2. Genome-wide association study implicates HLA-C*01:02 as a risk factor at the major histocompatibility complex locus in schizophrenia. *Biol Psychiatry.* 2012;72(8):620–628.
29. English JA, Lopez LM, O’Gorman A, et al. Blood-based protein changes in childhood are associated with increased risk for later psychotic disorder: evidence from a nested case-control study of the ALSPAC longitudinal birth cohort. *Schizophr Bull.* 2018;44(2):297–306.
30. Focking M, Sabherwal S, Cates HM, et al. Complement pathway changes at age 12 are associated with psychotic experiences at age 18 in a longitudinal population-based study: evidence for a role of stress. *Mol Psychiatry.* 2019;26(2):524–533.
31. Mongan D, Föcking M, Healy C, et al; European Network of National Schizophrenia Networks Studying Gene-Environment Interactions (EU-GEI) High Risk Study Group. Development of proteomic prediction models for transition to psychotic disorder in the clinical high-risk state and psychotic experiences in adolescence. *JAMA Psychiatry.* 2021;78(1):77–90.
32. Focking M, Pollak T, Dicker P, et al. Proteomic analysis of blood based samples from the optimise (optimization of treatment and management of schizophrenia in Europe) study point towards complement pathway protein changes. *Biol Psychiatry.* 2018;83(9):S301.
33. Leucht S, Winter-van Rossum I, Heres S, et al. The optimization of treatment and management of schizophrenia in Europe (OPTiMiSE) trial: rationale for its methodology and a review of the effectiveness of switching antipsychotics. *Schizophr Bull.* 2015;41(3):549–558.
34. Sheehan DV, Lecrubier Y, Sheehan KH, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry.* 1998;59(Suppl 20):22–33;quiz 34.
35. Andreasen NC, Carpenter WT, Jr, Kane JM, Lasser RA, Marder SR, Weinberger DR. Remission in schizophrenia: proposed criteria and rationale for consensus. *Am J Psychiatry.* 2005;162(3):441–449.
36. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull.* 1987;13(2):261–276. doi: [10.1093/schbul/13.2.261](https://doi.org/10.1093/schbul/13.2.261). PMID: 3616518.
37. Chiu E-C, Hung T-M, Huang C-M, Lee S-C, Hsieh C-L. Responsiveness of the personal and social performance scale in patients with schizophrenia. *Psychiatry Res.* 2018;260:338–342.
38. McGurk KA, Dagliati A, Chiasserini D, et al. The use of missing values in proteomic data-independent acquisition mass spectrometry to enable disease activity discrimination. *Bioinformatics.* 2020;36(7):2217–2223.
39. Rotello RJ, Veenstra TD. Mass spectrometry techniques: principles and practices for quantitative proteomics. *Curr Protein Pept Sci.* 2021;22(2):121–133.
40. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Methodol.* 1995;57(1):289–300.
41. Oikonomopoulou K, Ricklin D, Ward PA, Lambris JD. Interactions between coagulation and complement--their role in inflammation. *Semin Immunopathol.* 2012;34(1):151–165.
42. Tschopp J, Chonn A, Hertig S, French LE. Clusterin, the human apolipoprotein and complement inhibitor, binds to complement C7, C8 beta, and the b domain of C9. *J Immunol.* 1993;151(4):2159–2165.
43. Heurich M, Föcking M, Mongan D, Cagney G, Cotter DR. Dysregulation of complement and coagulation pathways: emerging mechanisms in the development of psychosis. *Mol Psychiatry.* 2021;27(1):127–140.
44. Enache D, Nikkheslat N, Fathalla D, et al. Peripheral immune markers and antipsychotic non-response in psychosis. *Schizophr Res.* 2021;230:1–8.
45. Martinuzzi E, Barbosa S, Daoudlarian D, et al; OPTiMiSE Study Group. Stratification and prediction of remission in first-episode psychosis patients: the OPTiMiSE cohort study. *Transl Psychiatry.* 2019;9(1):20.
46. Martinez-Cengotitabengoa M, MacDowell KS, Alberich S, et al; FLAMM-PEPs. BDNF and NGF signalling in early phases of psychosis: relationship with inflammation and response to antipsychotics after 1 year. *Schizophr Bull.* 2016;42(1):142–151.
47. Pollak TA, Vincent A, Iyegbe C, et al. Relationship between serum NMDA receptor antibodies and response to antipsychotic treatment in first-episode psychosis. *Biol Psychiatry.* 2021;90(1):9–15.
48. Tu C, Rudnick PA, Martinez MY, et al. Depletion of abundant plasma proteins and limitations of plasma proteomics. *J Proteome Res.* 2010;9(10):4982–4991.
49. Lee PY, Osman J, Low TY, Jamal R. Plasma/serum proteomics: depletion strategies for reducing high-abundance proteins for biomarker discovery. *Bioanalysis.* 2019;11(19):1799–1812.
50. Schubert OT, Gillet LC, Collins BC, et al. Building high-quality assay libraries for targeted analysis of SWATH MS data. *Nat Protoc.* 2015;10(3):426–441.