




REVIEW

Abandoning grassland management negatively influences plant but not bird or insect biodiversity in Europe

Tessa Elliott¹ | Amibeth Thompson¹  | Alexandra-Maria Klein¹  |
Christian Albert^{2,3} | Nico Eisenhauer^{4,5} | Florian Jansen⁶ |
Andrea Schneider⁷ | Martin Sommer⁸ | Tanja Straka⁹  | Josef Settele^{4,10} |
Maria Sporbert¹¹ | Franziska Tanneberger¹² | Anne-Christine Mupepele^{1,13}

¹Nature Conservation and Landscape Ecology, University of Freiburg, Freiburg, Germany

²Institute of Geography, Ruhr University Bochum, Bochum, Germany

³Institute of Environmental Planning, Leibniz University Hannover, Hannover, Germany

⁴German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany

⁵Institute of Biology, Leipzig University, Leipzig, Germany

⁶Landscape Ecology, University of Rostock, Rostock, Germany

⁷Faculty of Biology, Aquatic Ecology, University of Duisburg-Essen, Essen, Germany

⁸Deutscher Verband für Landschaftspflege (DVL)—Landcare Germany, Ansbach, Germany

⁹Institute of Ecology, Technische Universität Berlin, Berlin, Germany

¹⁰Helmholtz-Centre for Environmental Research—UFZ, Halle, Germany

¹¹Institute of Biology/Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle, Germany

¹²Greifswald Mire Centre, Greifswald University, Greifswald, Germany

¹³Department of Biology—Animal Ecology, University of Marburg, Marburg, Germany

Correspondence

Amibeth Thompson, Nature Conservation and Landscape Ecology, University of Freiburg, Tennenbacher Str. 4, 79106 Freiburg, Germany.

Email: amibeth.thompson@nature.uni-freiburg.de

Funding information

Deutsche Forschungsgemeinschaft, FZT 118, Grant/Award Number: 202548816; German Federal Ministry of Education and Research within the Research Initiative for the Conservation of Biodiversity (FEEdA), Grant/Award Number: 16LC2001B; Ministry of Science, Research and the Arts Baden-Württemberg

Abstract

Grasslands are globally distributed and naturally occurring; however, in Europe, most grasslands are anthropogenically created or altered by livestock grazing or mowing. Low-intensity use and management have led to species-rich communities in European grasslands. The intensification of crop production and livestock farming with stabling throughout the year has led to an abandonment of grasslands that are no longer economically profitable. In this study, we looked at the influence of grassland abandonment on biodiversity. We hypothesized that abandonment of grasslands decreases the overall biodiversity, but has different effects depending on the focal taxonomic group (i.e., vascular plants, insects, or birds). We also hypothesized that the type of management before abandonment, the type of grassland, and the time after abandonment would influence grassland biodiversity. We conducted a Web of Science search, with pre-defined terms, to find articles that compared biodiversity of managed and abandoned grasslands in Europe.

Tessa Elliott and Amibeth Thompson shared first authorship.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Conservation Science and Practice* published by Wiley Periodicals LLC on behalf of Society for Conservation Biology.

We screened the articles and included 39 studies in a subsequent meta-analysis. We found that overall biodiversity was reduced after abandonment; however, the biodiversity reduction in the grasslands differed among taxonomic groups. Plant species diversity was significantly lower after abandonment (plant summary effect size: -0.25 [-0.34 ; -0.16]), whereas the diversity of insects and birds showed no significant trend, but a visual trend toward an increase. None of the other environmental variables (type of management, type of grassland, or the time after abandonment) had a significant influence on the biodiversity of the grasslands. We conclude that maintaining grassland management is crucial to support biodiversity conservation in European grasslands.

KEYWORDS

bird, insect, meadow, meta-analysis, pasture, plant, species-richness

1 | INTRODUCTION

Grasslands are globally distributed and occur naturally if precipitation and temperature are within a given range (Dixon et al., 2014; Wilson et al., 2012; Woodward et al., 2004). In Europe, most grasslands have been anthropogenically used over centuries for agricultural production (Bardgett et al., 2021; Dengler et al., 2014; Feurdean et al., 2018; Pärtel et al., 2005; Pereira & Navarro, 2015; Vrahnakis et al., 2013). Grasslands are predominantly used for fodder production or direct livestock grazing (Hejman et al., 2013; Valkó et al., 2018; Wiezik et al., 2011). Low-intensity managed grasslands in Europe are not only important for the production of fodder, but can also contribute to biodiversity conservation (Dengler et al., 2014; Janišová et al., 2020). Along with anthropogenic grassland management, plant communities and diversity of higher trophic levels, such as herbivorous insects, have evolved to diverse systems (Cerabolini et al., 2016). Grasslands are habitats for many endemic or endangered plant species across different regions in Europe (Stoate et al., 2009). Supporting their conservation, numerous grassland types have been listed in the Annex I of the Habitats Directive (92/43/EEC), the main policy instrument for site protection at the EU level, such as mountain hay meadows and swards of Mat-grass (*Nardus stricta* L.; Evans & Roekarts, 2019; Henle et al., 2008).

Biodiversity in European grasslands is threatened most particularly by rapid and severe changes in agricultural production since the beginning of industrialization around 1830 (Habel et al., 2013; Mupepele et al., 2021; Queiroz et al., 2014). The invention of artificial fertilizer production has led to an intensification in grassland productivity, which favored herbaceous plant species, such as grasses. The additional nutrients from fertilizers reduce plant biodiversity and shift competition to other

limited resources such as light (Grace et al., 2016; Harpole et al., 2016). Livestock farming was equally industrialized and globalized with fodder production far beyond the local scale. The overall grassland area was reduced and fragmented by conversion to cropland or urban areas (Aune et al., 2018; Bengtsson et al., 2019; Pärtel et al., 2005; Schoof et al., 2019; Walcher et al., 2019).

Apart from the intensification and a general loss of grassland area, there is further a widespread trend of grassland abandonment in Europe, which has received less scientific attention (Henle et al., 2008; Nikolov, 2010). Abandonment occurs particularly on lands for which management no longer appears economically profitable, for instance due to difficult accessibility, trafficability, or rural population decline (Joyce, 2014; Valkó et al., 2018; Walcher et al., 2019). A consequence of management abandonment is a succession toward denser and taller vegetation (Tanneberger et al., 2008, 2010), and ultimately shrubby and forest vegetation (Bonanomi et al., 2013; Gibson, 2009; Shugart, 2013). Such an encroachment can in turn reduce grassland-associated species communities and their biodiversity (Azcárate & Peco, 2012; Facioni et al., 2019; Pärtel et al., 2005; Pereira & Navarro, 2015). In other cases, management abandonment led to habitat regeneration and increased biodiversity (Azcárate & Peco, 2012; Kuhn et al., 2021; Laiolo et al., 2004; Skórka et al., 2007). Thus, succession can be progressive, leading to increased biodiversity, structural complexity, and stability, or retrogressive, changing an ecosystem in the opposite direction and reducing biodiversity (Shugart, 2013). Abandonment has been known to both decrease (Azcárate & Peco, 2012; Pärtel et al., 2005) and increase biodiversity (Azcárate & Peco, 2012), but a comprehensive quantitative synthesis of these findings and their difference between management, time since abandonment, or habitat characteristics is still missing.

Based on the previous findings that management, such as mowing and subsequent biomass removal in low intensities, is a key aspect for maintaining biodiversity in grasslands (Cerabolini et al., 2016; Dengler et al., 2014; Valkó et al., 2018), we hypothesized that overall, abandonment of grasslands decreases the biodiversity and has differential effects depending on the taxonomic group studied. We also expected that the type of management, the type of grassland, and the years after abandonment would influence the biodiversity of grasslands. To identify an overall change in biodiversity potentially influenced by environmental variables, we synthesized the existing information of biodiversity on abandoned grasslands in Europe in the form of a meta-analysis to address the following research questions:

1. Does the abandonment of grassland in Europe have an impact on overall biodiversity?
2. Does the effect of abandonment on biodiversity differ among taxonomic groups?
3. What other factors, such as the type of previous management, for example, mowing or grazing, the type of grassland ranging from dry to moist, or the time after abandonment, influence biodiversity changes in abandoned European grasslands?

2 | METHODS

2.1 | Literature search

Studies published between 1995 and 2021 comparing biodiversity on managed and abandoned grasslands in Europe were searched in Clarivate Web Of Science (WOS, <https://www.webofscience.com>) in July 2021. We used search terms related to “biodiversity,” “grassland,” “abandonment,” and “Europe” (full search string in Appendix S1), which resulted in 1079 potentially relevant studies. Studies revealed by the WOS search had to fulfill additional inclusion criteria, namely the study (i) was conducted in Europe, (ii) looked at the biodiversity of a grassland (grassland definition based on Dengler et al., 2014), and (iii) compared abandoned with managed (control) grassland sites (full list of criteria in Table 1). The 1079 potentially relevant articles resulting from the WOS search were screened to identify whether they fulfilled all inclusion criteria. According to general systematic review standards, we first screened title and abstract and discarded studies not fulfilling the inclusion criteria (e.g., if the abstract revealed that the study was conducted outside of Europe). In a second screening, the full-text was read and studies not fulfilling the inclusion criteria were again discarded. After the screenings, 39 studies

TABLE 1 Criteria used for studies to be included or excluded in this review.

Criteria	Eligibility	Exclusion
Study location	Study sites located within Europe	Study sites located outside of Iceland to the Ural Mountains and from Norway to the Mediterranean Sea and the Black Sea
	Study was conducted in a grassland which means densely-covered herbaceous vegetation dominated by grasses (Dengler et al., 2014)	Study was conducted in a different type of habitat or in a grassland not as defined
Language	In English or German	Other languages outside of English or German
Accessibility	Article accessible through the subscriptions held by the University of Freiburg or personal communication with the authors	Articles inaccessible through the subscription or through personal communication
Study design	Study recorded data on a grassland on which management had been abandoned as well as on a control site with management in the form of biomass removal	Sites on which management had recently changed, for example, by adding irrigation or fertilizer, or which were restored after a period of abandonment
	The management on the abandoned grassland and control grassland are clearly described and replicable	The differentiation of abandoned and managed grassland based solely on visual characteristics such as “degree of shrub encroachment”
	Study measured average species richness or other quantifiable biodiversity measures, such as Shannon diversity, and data are	Study did not measure biodiversity or the measure was not extractable

(Continues)

TABLE 1 (Continued)

Criteria	Eligibility	Exclusion
	extractable for an abandoned grassland site and a corresponding managed control grassland in relation to their sample size and a variance measure	

were left that met all inclusion criteria (see Table 1, Appendix S2). Environmental variables (type of management, type of grassland, or the time after abandonment) and the biodiversity metrics (species richness [preferential], Shannon-Diversity-Index, species density) with sample size and standard deviation, were extracted, each for an abandoned grassland (treatment) and a managed grassland (control). Unique control–treatment combinations of control types, taxonomic groups, and study site locations were extracted, which meant that some studies provided several combinations. We extracted a total of 90 control–treatment combinations, that is, 90 effect sizes, out of the 39 studies, and, if possible, their concordant environmental variables (see Appendix S3 and S4). WebPlotDigitizer was used to extract data points from figures (Rohatgi, 2021). If standard deviation was not given, but other variance measures, such as standard error, they were converted to standard deviation (Higgins et al., 2021). If no information about variance or sample size was given and the study was published within the last 10 years, we contacted the authors via e-mail and asked for further information that would allow us to compute the standard deviation. Some studies used a nested design with several data points on a larger plot so that information about the diversity on a small scale plot level, for example, 1 m², and a larger scale site level, for example, 1 km² were available. Only a few studies (9 from 39) provided information on the size of the area, so that the influence of the spatial scale could not be included in the analysis. If diversity was given on several scales, only the largest scale was included in the analysis. For studies that provided data over several years after the abandonment, only the last data point in a time series was used to guarantee an equal weighting for all study sites across the meta-analysis.

2.2 | Dataset description

The final dataset is composed of 90 effect sizes from studies published between 1995 and 2021 with data from

19 countries in Europe (see Appendix S5). We classified taxonomic groups into vascular plants (64 effect sizes), insects (18 effect sizes), and birds (8 effect sizes) as there were no effect sizes with other invertebrates and vertebrates. The orders of insects that were studied were granivorous Formicidae (ants), phytophagous, saprophagous, and coprophagous Coleoptera (beetles), and diurnal Lepidoptera (butterflies). Grassland management type was well balanced with 52 effect sizes for grazed grasslands, 31 effect sizes for mowed grasslands, and a further 6 effect sizes with grassland being grazed and mowed. One effect size with a control site subject to controlled burning was excluded from the mixed-effects model analysis moderated by grassland management (see below). The grasslands were categorized into three different types: dry grasslands (45 effect sizes), mesic grasslands (14 effect sizes), or wet to seasonally wet grasslands (7 effect sizes). Only 73% of the studies reported the type of grassland and 68% of the studies focused on dry grasslands. About 74% of the studies reported the years since abandonment, which ranged from 2 to 60 years, with 15.4 as the average (Appendix S7). Our results are dominated by plant studies (64 out of 90 effect sizes), dry grasslands (45 out of 66 effect sizes), and grasslands abandoned less than 20 years ago (47 out of 67 effect sizes).

2.3 | Statistical analysis

The log response ratio was chosen as an effect size to compare the biodiversity of abandoned grasslands (“treatment”) with grasslands managed, for example, by mowing or grazing (“control”) in a meta-analysis (Gurevitch et al., 2018; Koricheva et al., 2013). A log response ratio below zero reflects a decrease in biodiversity after management abandonment. The log response ratio was chosen as we were interested in the proportional change of biodiversity, reflected in ratios (Hillebrand & Gurevitch, 2016). Log response ratios can further be used if measurement units differ between studies, such as in our case, in which we had effect sizes derived from species richness or Shannon diversity. The summary effect size was estimated based on a three-level random-effects model (Konstantopoulos, 2011; Nakagawa & Santos, 2012). This model was chosen to incorporate the hierarchies and dependencies in the dataset resulting from the fact that some effect sizes originated from the same study. A three-level random-effect model assumes three sources of variance with sampling error as well as two sources of heterogeneity (Konstantopoulos, 2011; Nakagawa & Santos, 2012). To identify whether there was a high amount of variance attributed to between-study variability (heterogeneity) or within-study variability two approaches were used: (i) a Q-test, which can be seen as an

ANOVA counterpart in meta-analysis and reflects the distance of effect sizes from primary studies to the mean effects size, and (ii) the index I^2 , specifying the proportion of the between-study variability in relation to the total variance, was used (Assink & Wibbelink, 2016). To consider variables possibly explaining the between-study variability, we used mixed-effects models and a subsequent omnibus test to identify a potential moderating effect of the environmental variables “taxonomic group,” “management on the control grassland,” “grassland type,” and “time since abandonment” (Table 2). The moderating effect of the four environmental variables was assessed in four different models, as not all studies accounted for all environmental variables simultaneously, and a different subset of studies was included in each mixed-effects model analysis. To account for multiple testing, a Bonferroni correction was employed, leading to a modified significance level ($P = \alpha/4 = 0.0125$; Nakagawa, 2004). If the omnibus test revealed an influence of one of the categorical environmental variables (i.e., “taxonomic group,” “management on the control grassland,” or “grassland type”), a subsequent sub-group analysis was realized, which can be seen as the counterpart to a post-hoc test providing the information for which of the categories

TABLE 2 Mixed-effects models to investigate whether an environmental variable influences the effect sizes, that is, the change in biodiversity on grasslands after abandonment.

Model identifier	Environmental variable	Categories and units of environmental variables
Model 1	Taxonomic group	Vegetation (vascular plants), invertebrates (insects), vertebrates (birds)
Model 2	Grassland management	Mown, grazed, mixed management, controlled fire
Model 3	Grassland type	Dry grassland, mesic grassland, seasonally wet and wet grassland (for manual classification additional literature was consulted: Evans & Roekarts, 2019; Sauermost et al., 1999; Spohn et al., 2015)
Model 4	Time since abandonment	Number of years (if a period was given, the average was used; in case of abandonment of more than a specified period of time the given value was adopted [e.g., >10 years became 10 years])

biodiversity significantly increased or decreased after abandonment (Rossetti et al., 2017).

We assessed the robustness and generalizability of the statistical results, in relation to a potential publication bias (Gurevitch & Nakagawa, 2015). A publication bias can occur if the effect sizes in a meta-analysis are biased, for example, toward significant results or toward supporting only a certain narrative. The publication bias is also referred to as a “file-drawer-problem” alluding to non-significant results that are hidden in the drawer of researchers instead of being published. They are thus not found in a literature search and missing in a meta-analysis, which is biasing the results (Koricheva et al., 2013). We investigated the presence of publication bias visually based on a funnel plot (Sterne & Harbord, 2004; Sterne et al., 2011) and by means of Egger's regression asymmetry test (Egger et al., 1997; Koricheva et al., 2013; adapted from Mupepele et al., 2021). Analyses were conducted in R 4.0.5 using the package “metafor” (Viechtbauer, 2010, see Appendix S9).

3 | RESULTS

Biodiversity was reduced in abandoned compared to managed grasslands (summary effect size: -0.14 [-0.24 ; -0.04], Figure 1, Appendix S8). Only a small proportion of variance was linked to sampling variance (I^2 [level 1] = 1.09%). A high variability exists among studies demonstrating the importance of investigating environmental variables potentially explaining parts of this variance (between-study variance: Log-likelihood ratio test [LRT] = 18.79; $P < 0.0001$; I^2 [level 3] = 57.54%; within-study variance: LRT = 1858.4; $P < 0.0001$; I^2 [level 2] = 41.37%). Neither grassland management, such as mowing or grazing, nor the grassland type, such as dry or wet grassland, nor time that the grassland had been abandoned had an impact on the change in biodiversity after abandonment (Table 3, Appendix S7). There was, however, a significant difference between the taxonomic groups (Table 3, Model 1). Plant diversity decreased after abandonment (Figure 1, green diamond, Plant summary effect size: -0.25 [-0.34 ; -0.16]). Insects and birds did not show statistical evidence for a change in biodiversity, but a visual trend toward an increasing biodiversity.

3.1 | Publication bias

According to the funnel plot and Egger's regression test, no publication bias was detected in our data (Appendix S6; intercept of Egger's regression = -0.82 ; $t = -0.65$; $P = 0.52$).




FIGURE 1 Forest plot demonstrating an overall decrease in biodiversity after abandoning grassland management (black diamond represents the summary effect size). The overall decrease in biodiversity was dominated by a reduced plant diversity (green diamond represents the subgroup summary effect size for plants), while insects (yellow) and birds (blue) were showing a non-significant trend toward an increase in diversity. Effect sizes of primary studies are represented by a gray point and their corresponding confidence intervals.

TABLE 3 Mixed-effects model results for each environmental variable potentially influencing the effect sizes.

Model number	Environmental variable	Number of effect sizes ^a	Omnibus-test, Q_M (df)	P-value
1	Taxonomic group	90	60.35 (3)	<0.0001*
2	Grassland management	89	9.12 (3)	0.0277 ^b
3	Grassland type	66	3.2505 (3)	0.36
4	Time since abandonment	67	1.25 (1)	0.26

^aEffect sizes represent differences in the biodiversity of managed and abandoned grasslands.

^bStatistically significant according to the standard level of significance ($P < 0.05$), non-significant after the Bonferroni correction.

*Statistically significant according to the modified level of significance after the Bonferroni correction ($P < 0.0125$).

4 | DISCUSSION

Our meta-analysis synthesized findings across 39 studies with 90 effect sizes comparing managed and abandoned grasslands and demonstrated that biodiversity decreases after abandoning grassland management in Europe, which was driven by a loss in plant diversity. Insect and bird diversity showed a positive trend, which could indicate a trophic cascading effect, but additional studies would be needed to support this currently non-significant trend. Contrary to our hypotheses, we found no evidence for an influence of the type of management, such as mowing or grazing, the type of grassland, and the time since abandonment on the change in biodiversity.

4.1 | Plant diversity in abandoned grasslands

Abandoning grassland management leads to succession in plant communities and can thus reduce plant species diversity (Shugart, 2013; Walcher et al., 2019). Due to the

missing disturbances, such as animal trampling and biomass removal by mowing or grazing, plant litter accumulates. Plant litter can be beneficial for plants by conserving water in dry periods, but the overall effects on vegetation communities are negative (Pärtel et al., 2005; Valkó et al., 2012; Xiong & Nilsson, 1999). Litter may alter the physical and chemical environmental factors of the upper soil layer, which can reduce germination rates of certain species (Valkó et al., 2012). Additionally, the higher availability of nutrients contained in the litter causes eutrophication and increased productivity and canopy height. This is shifting resource competition of plants toward the availability of light (Hautier et al., 2009; Joyce, 2014) and thus, few fast-growing species might outcompete less dominant species, which are progressively replaced (Dierschke & Briemle, 2002; Hautier et al., 2009; Joyce, 2014; Otsus et al., 2014). With ongoing succession, the average vegetation height increases, and shrub encroachment as well as woody vegetation can inhibit the growth of short herbaceous species due to a lack of disturbance and light (Pápay et al., 2020; Pöyry et al., 2006). Our results found that abandonment

decreases plant biodiversity and that management, such as mowing and subsequent biomass removal as a regular disturbance, is important for a positive response of plant diversity (Dengler et al., 2014; Pöyry et al., 2006; Tanneberger et al., 2022; Xiong & Nilsson, 1999). The general decrease of plant species after grassland abandonment does not exclude the existence of plant species specialized and dependent on abandonment or the process of succession and we are recognizing that the biology of certain species (i.e., grasses vs. herbs species) can strongly affect the reaction to a specific management (i.e., grazing vs. mowing, Steffan-Dewenter & Leschke, 2003). While the majority of case studies in our meta-analysis were focused on plants, the knowledge on grassland fauna is more limited (Dengler et al., 2014; Joyce, 2014; Walcher et al., 2019).

4.2 | Insect diversity in abandoned grasslands

We found no significant effect of management abandonment on insects. The studies vary in the focal groups, such as ants, beetles, or butterflies, and thus they also vary in their traits, including their mobility, body size, and feeding and nesting needs. These differences can lead to different responses to management and its abandonment. For example, herbivorous insects directly depend on plants; however, they may react with a delay to changes in the lower trophic level, such as plant diversity. Three of the seven included studies looked at functional groups or traits of the insects, with only one study specifically looking at herbivorous insects (Fadda et al., 2008). The sample size was too low to further split insects into lower taxonomic or functional groups during the analysis. Likewise, the effects could differ between management types, with grazing having a different effect in comparison to mowing (Bonari et al., 2017) or between habitat types, with wet meadows having a different effect from dry meadows (Skórka et al., 2007). The lack of management could locally increase habitat heterogeneity with potentially positive effects for the insect population (Azcárate & Peco, 2012; Laiolo et al., 2004; WallisDeVries et al., 2002). Differences within species groups (i.e., large vs. small body size) also show diverging reactions to changes in the grasslands (Tonelli et al., 2018). Generalist species are usually more robust to changes and thus possibly less sensitive to management abandonment (Öckinger et al., 2006). In our analysis, only one study looked explicitly at specialist species and found that habitat type was the main driver and not management (Wiezik et al., 2019). Likewise, only four of the studies looked at the plant community in conjunction with

the insect group, making it impossible to draw any conclusions on the influences of grassland abandonment on the overall insect communities. The decrease in disturbance is also known to explain an increase in abundance and diversity in arthropod communities (Morris, 2000), especially for insects that are also strongly related to soil (i.e., soil larvae, dormancy), which may be promoted by the lack of disturbance and accumulated litter during abandonment.

4.3 | Bird diversity in abandoned grasslands

Similar to insects, we did not find significant effects of abandonment of grassland management for birds, which could be explained by the fact that abandonment initially has a positive effect on the diversity of fauna, and insects and birds may initially benefit from the increased structural complexity (Verholst et al., 2004). For example, less disturbance creates greater vegetation height for cover or more flowers for pollinators (Laiolo et al., 2004). However, if a site is no longer managed and in the long term, a thick layer of litter accumulates (Tanneberger et al., 2008, 2010) or the site becomes woody, grassland-dependent insect species will decline and, in turn, the birds that feed on those insects may decline (Kobbeloer & Lanz, 2018). On the one hand, birds have a larger habitat range than plants and insects, and thus the change or lack of management may not have any short-term implications on their community. On the other hand, insects depend on plants as a food source, and plants depend on insects for pollination. Likewise, birds rely on insects as food source and ground-nesting birds also on vegetation structure for nesting and cover (Arbeiter et al., 2018, 2020). Since each of these groups are interdependent on one another, we predicted that a significant decrease in plant species would also ultimately lead to a decrease of insect and bird species, which we could not confirm.

4.4 | Influence of time since abandonment, grassland type, and management type

We expected that ongoing succession, and thus the time since abandonment, would have an influence on the difference of biodiversity between managed and abandoned grasslands. We could not confirm this effect; eventually including and accounting for the progression within long-term studies may shed more light on this relationship. While there are scientific records of short periods of abandonment promoting certain species and leading to

an increase in biodiversity, long-term abandonment often leads to population decline and local extinction events (Barabasz-Krasny, 2011; Bonari et al., 2017; Valkó et al., 2012). For instance, abandoned grasslands can still serve as a habitat for insects specialized on open land until woody vegetation encroaches (Wiezik et al., 2011). The time in which negative effects of abandonment arise have been shown to differ between different plant communities, depending on factors such as productivity (Bonari et al., 2017). Thus, the disruption of biodiversity seems to occur at different states of successions, which can be stable for several years to decades depending on the stage, abiotic and biotic factors and other variables such as management history (Walcher et al., 2019; Wiezik et al., 2011). Since succession can be fast or slow depending on the local conditions (Joyce, 2014), studies defining successional stages in addition to the number of years since management abandonment may support conclusions in biodiversity changes.

We also expected that different grassland types would influence biodiversity. Different types usually have different species composition and thus responses to environmental conditions and changes (e.g., Evans & Roekarts, 2019). However, we could not confirm a significant difference between grassland types in the biodiversity change after abandonment. In a previous meta-analysis, it was found that specialist plant species in wet grasslands almost disappear completely over time, whereas it was not so dramatic for dry grasslands (Diekmann et al., 2019). Likewise, differences between fauna composition have been found between the different types of grasslands, specifically for butterflies (Öckinger et al., 2006; Skórka et al., 2007), with abandonment of wet meadows having a greater, positive impact on the community. Eventually, a more equal distribution of effect sizes across the different grassland types could reveal differences between dry, mesic, and wet grasslands in biodiversity change after abandonment.

Different management practices are known to lead to different species communities in grasslands, especially for plants (Dierschke & Briemle, 2002; Kahmen et al., 2002; Kuhn et al., 2021; Rudmann-Maurer et al., 2008), but the effects on biodiversity are less apparent (Dengler et al., 2014). We expected that the management which had taken place before abandonment would have influenced the change in diversity. Grazing in comparison to mowing was found to promote higher levels of biodiversity, particularly for butterflies and ground beetles (Tälle et al., 2016). Abandonment of grazed sites could suffer from a loss of small-scale heterogeneity that was caused by trampling and defecation patterns (Dengler et al., 2014; Tälle et al., 2016). In our analysis, we did not find any significant influence of the management type taking place

before abandonment, that is, grazing, mowing, or both. On the one hand, a possible explanation is that spatial context of the landscape is important with the difference between local and regional scale or the complexity of the landscape shaping the diversity of different communities (Gossner et al., 2016; Le Provost et al., 2021). On the other hand, temporal stability of management might be of higher importance for biodiversity than the type of management in grasslands (Eriksson et al., 2002). Likewise, historical management can play an important role for the established species community as well as the suitability of the present land-use (Tälle et al., 2016). Management as well as its abandonment also has an important impact on the soil community; however, current nature protection does not seem to benefit soil biodiversity (Ciobanu et al., 2019) or functions (Zeiss et al., 2022). Therefore, more tailored management and conservation approaches are needed not just for the aboveground communities, but also for the belowground communities (Guerra et al., 2022).

4.5 | Conservation and future research

Grassland management has a long-standing tradition in European landscapes. Not only is there a diversity of grassland types, but grasslands are essential in promoting biodiversity and ecosystem services (Schils et al., 2022). In our analysis, we demonstrate that abandonment reduces biodiversity, most particularly plant species richness. Beyond plants, we encourage future comparative studies (using control sites) for insects (also including below ground species) and birds as our data did not yet reveal a clear understanding of how their diversity is affected by grassland management abandonment. We were further surprised that our literature search revealed missing studies in taxonomic groups other than plants, insects, and birds and the lack of studies studying the effects of specialists versus generalists. The industrialization of agriculture has changed grassland management as well as its diversity, involving both intensification in productive areas and abandonment in economically less attractive areas. Due to the previously non-uniform ways to report management intensity (e.g., “number of grazers/ha” in Jacquemyn et al., 2011; “estimated grazing pressure” in Laiolo et al., 2004), we were unable to include the management intensity on the control sites as a variable in our meta-analysis. Midolo et al. (2023) have suggested a uniform way of reporting disturbance indicator values, which would allow quantifying disturbance and intensification across different grasslands and studies. The bias toward plants, nonuniform methods, lack of a control, or incomplete data continue to be issues when

conducting a meta-analysis on grasslands and management (Slodowicz et al., 2023) and should be considered for future grassland studies. Considering different biodiversity facets or the spatial context of the landscape may also provide useful insight into understanding the biodiversity of the different communities. Given the overall decline in biodiversity after abandoning grassland management, we further encourage policy measures to support farmers maintaining their traditional, extensively managed grasslands.

ACKNOWLEDGMENTS

We would like to thank all authors of the primary publications included in this meta-analysis for their support and the additional data and information they eventually provided. Open Access funding enabled and organized by Projekt DEAL.

FUNDING INFORMATION

This project was funded by the German Federal Ministry of Education and Research within the Research Initiative for the Conservation of Biodiversity (FEa) under the funding code 16LC2001B. The responsibility for the content of this publication lies with the authors. Nico Eisenhauer acknowledges support of iDiv funded by the German Research Foundation (DFG-FZT 118, 202548816). Anne-Christine Mupepele was supported by the Ministry of Science, Research and the Arts Baden-Württemberg, Germany.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article (Appendix S9 and S10).

ORCID

Amibeth Thompson  <https://orcid.org/0000-0002-9224-412X>

Alexandra-Maria Klein  <https://orcid.org/0000-0003-2139-8575>

Tanja Straka  <https://orcid.org/0000-0003-4118-4056>

REFERENCES

- Arbeiter, S., Flinks, H., Grünwald, J., & Tanneberger, F. (2020). Diet of corncrakes *Crex crex* and prey availability in relation to meadow management. *Ardea*, 108(1), 55–64.
- Arbeiter, S., Roth, T., Helmecke, A., Haferland, H. J., Tanneberger, F., & Bellebaum, J. (2018). Conflict between habitat conservation and corncrake *Crex crex* brood protection in managed floodplain meadows. *Agriculture, Ecosystems & Environment*, 265, 15–21.
- Assink, M., & Wibbelink, C. J. (2016). Fitting three-level meta-analytic models in R: A step-by-step tutorial. *The Quantitative Methods for Psychology*, 12, 154–174.
- Aune, S., Bryn, A., & Hovstad, K. A. (2018). Loss of semi-natural grassland in a boreal landscape: Impacts of agricultural intensification and abandonment. *Journal of Land Use Science*, 13, 375–390.
- Azcárate, F. M., & Peco, B. (2012). Abandonment of grazing in a mediterranean grassland area: Consequences for ant assemblages. *Insect Conservation and Diversity*, 5, 279–288.
- Barabasz-Krasny, B. (2011). Vegetation dynamics in fallow agricultural areas on Przemyśl foothills (southeastern Poland). *Acta Societatis Botanicorum Poloniae*, 74, 149–157.
- Bardgett, R. D., Bullock, J. M., Lavorel, S., Manning, P., Schaffner, U., Ostle, N., & Bullock, J. M. (2021). Combating global grassland degradation. *Nature Reviews and Earth Environment*, 2, 720–735.
- Bengtsson, J., Bullock, J. M., Egoh, B., Everson, C., Everson, T., O'Connor, T., & Bullock, J. M. (2019). Grasslands-more important for ecosystem services than you might think. *Ecosphere*, 10, e02582.
- Bonanomi, G., Incerti, G., & Allegranza, M. (2013). Assessing the impact of land abandonment, nitrogen enrichment and fairy-ring fungi on plant diversity of Mediterranean grasslands. *Biodiversity and Conservation*, 22, 2285–2304.
- Bonari, G., Fajmon, K., Malenovský, I., Zelený, D., Holuša, J., Jongepierová, I., & Fajmon, K. (2017). Management of semi-natural grasslands benefiting both plant and insect diversity: The importance of heterogeneity and tradition. *Agriculture, Ecosystems & Environment*, 246, 243–252.
- Cerabolini, B. E. L., Pierce, S., Verginella, A., Brusa, G., Ceriani, R. M., & Armiraglio, S. (2016). Why are many anthropogenic agroecosystems particularly species-rich? *Plant Biosystems—An International Journal Dealing with all Aspects of Plant Biology*, 150, 550–557.
- Ciobanu, M., Eisenhauer, N., Stoica, I.-A., & Cesarz, S. (2019). Natura 2000 priority and non-priority habitats do not differ in soil nematode diversity. *Applied Soil Ecology*, 135, 166–173.
- Dengler, J., Janišová, M., Török, P., & Wellstein, C. (2014). Biodiversity of Palearctic grasslands: A synthesis. *Agriculture, Ecosystems & Environment*, 182, 1–14.
- Diekmann, M., Andres, C., Becker, T., Bennie, J., Blüml, V., Bullock, J. M., & Andres, C. (2019). Patterns of long-term vegetation change vary between different types of semi-natural grasslands in Western and Central Europe. *Journal of Vegetation Science*, 30, 187–202.
- Dierschke, H., & Briemle, G. (2002). *Kulturgrasland: Wiesen, Weiden und verwandte Staudenfluren ; 20 Tabellen*. Ökosysteme Mitteleuropas aus geobotanischer Sicht. Ulmer, Stuttgart.
- Dixon, A. P., Faber-Langendoen, D., Josse, C., Morrison, J., & Loucks, C. J. (2014). Distribution mapping of world grassland types. *Journal of Biogeography*, 41, 2003–2019.
- Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315, 629–634.
- Eriksson, O., Cousins, S. A. O., & Bruun, H. H. (2002). Land-use history and fragmentation of traditionally managed grasslands in Scandinavia. *Journal of Vegetation Science*, 13, 743–748.
- Evans, D., & Roekarts, M. (2019). Interpretation manual of the habitats listed in resolution No. 4 (1996) listing endangered natural habitats requiring specific conservation measures.
- Facioni, L., Burrascano, S., Chiti, T., Giarrizzo, E., Zanini, M., & Blasi, C. (2019). Changes in plant diversity and carbon stocks along a succession from semi-natural grassland to submediterranean

- Quercus cerris* L. Woodland in Central Italy. *Phytocoenologia*, *49*, 393–408.
- Fadda, S., Henry, F., Orgeas, J., Ponel, P., Buisson, E., & Dutoit, T. (2008). Consequences of the cessation of 3000 years of grazing on dry Mediterranean grassland ground-active beetle assemblages. *Comptes Rendus Biologies*, *331*, 532–546.
- Feurdean, A., Ruprecht, E., Molnár, Z., Hutchinson, S. M., & Hickler, T. (2018). Biodiversity-rich European grasslands: Ancient, forgotten ecosystems. *Biological Conservation*, *228*, 224–232. <https://doi.org/10.1016/j.biocon.2018.09.022>
- Gibson, D. J. (2009). *Grasses and grassland ecology*. Oxford University Press.
- Gossner, M. M., Lewinsohn, T. M., Kahl, T., Grassein, F., Boch, S., Prati, D., & Lewinsohn, T. M. (2016). Land-use intensification causes multitrophic homogenization of grassland communities. *Nature*, *540*, 266–269.
- Grace, J. B., Anderson, T. M., Seabloom, E. W., Borer, E. T., Adler, P. B., Harpole, W. S., & Anderson, T. M. (2016). Integrative modelling reveals mechanisms linking productivity and plant species richness. *Nature*, *529*, 390–393.
- Guerra, C. A., Berdugo, M., Eldridge, D. J., Eisenhauer, N., Singh, B. K., Cui, H., & Berdugo, M. (2022). Global hotspots for soil nature conservation. *Nature*, *1–6*, 610.
- Gurevitch, J., Koricheva, J., Nakagawa, S., & Stewart, G. (2018). Meta-analysis and the science of research synthesis. *Nature*, *555*, 175–182.
- Gurevitch, J., & Nakagawa, S. (2015). Chapter 9: Research synthesis methods in ecology. In G. A. Fox, S. Negrete-Yankelevich, & V. J. Sosa (Eds.), *Ecological statistics: Contemporary theory and application*. Oxford University Press.
- Habel, J. C., Dengler, J., Janišová, M., Török, P., Wellstein, C., & Wiezik, M. (2013). European grassland ecosystems: Threatened hotspots of biodiversity. *Biodiversity and Conservation*, *22*, 2131–2138.
- Harpole, W. S., Sullivan, L. L., Lind, E. M., Firn, J., Adler, P. B., Borer, E. T., & Sullivan, L. L. (2016). Addition of multiple limiting resources reduces grassland diversity. *Nature*, *537*, 93–96.
- Hautier, Y., Niklaus, P. A., & Hector, A. (2009). Competition for light causes plant biodiversity loss after eutrophication. *Science*, *324*, 636–638.
- Hejcman, M., Hejcmanová, P., Pavlů, V., & Beneš, J. (2013). Origin and history of grasslands in Central Europe—a review. *Grass and Forage Science*, *68*, 345–363.
- Henle, K., Alard, D., Clitherow, J., Cobb, P., Firbank, L., Kull, T., & Alard, D. (2008). Identifying and managing the conflicts between agriculture and biodiversity conservation in Europe—a review. *Agriculture, Ecosystems & Environment*, *124*, 60–71.
- Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., & Welch, V. (2021). *Cochrane handbook for systematic reviews of interventions: Version 6.2*. <https://training.cochrane.org/handbook/current>
- Hillebrand, H., & Gurevitch, J. (2016). Meta-analysis and systematic reviews in ecology. In *eLS* (pp. 1–11). John Wiley & Sons, Ltd.
- Jacquemyn, H., Mechelen, C. V., Brys, R., & Honnay, O. (2011). Management effects on the vegetation and soil seed bank of calcareous grasslands: An 11-year experiment. *Biological Conservation*, *144*, 416–422.
- Janišová, M., Biro, A., Iuga, A., Šírka, P., & Škodová, I. (2020). Species-rich grasslands of the Apuseni Mts (Romania): Role of traditional farming and local ecological knowledge. *Tuexenia*, *40*, 409–427.
- Joyce, C. B. (2014). Ecological consequences and restoration potential of abandoned wet grasslands. *Ecological Engineering*, *66*, 91–102.
- Kahmen, S., Poschlod, P., & Schreiber, K.-F. (2002). Conservation management of calcareous grasslands. Changes in plant species composition and response of functional traits during 25 years. *Biological Conservation*, *104*, 319–328.
- Kobbeloer, D., & Lanz, U. (2018). *Naturschutz Und Landwirtschaft Hand in Hand: das Bayerische Artenhilfsprogramm für den Ortolan*, 9.
- Konstantopoulos, S. (2011). Fixed effects and variance components estimation in three-level meta-analysis. *Research Synthesis Methods*, *2*, 61–76.
- Koricheva, J., Gurevitch, J., & Mengersen, K. (2013). *Handbook of meta-analysis in ecology and evolution*. Princeton University Press.
- Kuhn, T., Domokos, P., Kiss, R., & Ruprecht, E. (2021). Grassland management and land use history shape species composition and diversity in Transylvanian semi-natural grasslands. *Applied Vegetation Science*, *24*, e12585.
- Laiolo, P., Dondero, F., Ciliento, E., & Rolando, A. (2004). Consequences of pastoral abandonment for the structure and diversity of the alpine avifauna. *Journal of Applied Ecology*, *41*, 294–304.
- Le Provost, G., Thiele, J., Westphal, C., Penone, C., Allan, E., Nyret, M., van der Plas, F., Ayasse, M., Bardgett, R. D., Birkhofer, K., Boch, S., Bonkowski, M., Buscot, F., Feldhaar, H., Gaulton, R., Goldmann, K., Gossner, M. M., Klaus, V. H., Kleinebecker, T., ... Manning, H. (2021). Contrasting responses of above- and belowground diversity to multiple components of land-use intensity. *Nature Communications*, *12*, 3918.
- Midolo, G., Herben, T., Axmanová, I., Marcenò, C., Pätzsch, R., Bruelheide, H., Karger, D. N., Acíć, S., Bergamini, A., Bergmeier, E., Biurrun, I., Bonari, G., Čarni, A., Chiarucci, A., de Sanctis, M., Demina, O., Dengler, J., Dziuba, T., Fanelli, G., ... Chytrý, M. (2023). Disturbance indicator values for European plants. *Global Ecology and Biogeography*, *32*, 24–34.
- Morris, M. G. (2000). The effects of structure and its dynamics on the ecology and conservation of arthropods in British grasslands. *Biological Conservation*, *95*, 129–142.
- Mupepele, A.-C., Bruelheide, H., Brühl, C., Dauber, J., Fenske, M., Freibauer, A., & Bruelheide, H. (2021). Biodiversity in European agricultural landscapes: Transformative societal changes needed. *Trends in Ecology & Evolution*, *36*, 1067–1070.
- Mupepele, A.-C., Keller, M., & Dormann, C. F. (2021). European agroforestry has no unequivocal effect on biodiversity: A time-cumulative meta-analysis. *BMC Ecology and Evolution*, *21*, 193.
- Nakagawa, S. (2004). A farewell to Bonferroni: The problems of low statistical power and publication bias. *Behavioral Ecology*, *15*, 1044–1045.
- Nakagawa, S., & Santos, E. S. (2012). Methodological issues and advances in biological meta-analysis. *Evolutionary Ecology*, *26*, 1253–1274.
- Nikolov, S. C. (2010). Effects of land abandonment and changing habitat structure on avian assemblages in upland pastures of Bulgaria. *Bird Conservation International*, *20*, 200–213.

- Öckinger, E., Eriksson, A. K., & Smith, H. G. (2006). Effects of grassland abandonment, restoration and management on butterflies and vascular plants. *Biological Conservation*, *133*, 291–300.
- Otsus, M., Kukkk, D., Kattai, K., & Sammul, M. (2014). Clonal ability, height and growth form explain species' response to habitat deterioration in Fennoscandian wooded meadows. *Plant Ecology*, *215*, 953–962.
- Pápay, G., Kiss, O., Fehér, Á., Szabó, G., Zimmermann, Z., Hufnagel, L., Járđi, I., Szemethy, L., Penksza, K., & Katona, K. (2020). Impact of shrub cover and wild ungulate browsing on the vegetation of restored mountain hay meadows. *Tuexenia*, *40*, 445–457.
- Pärtel, M., Bruun, H. H., & Sammul, M. (2005). Biodiversity in temperate European grasslands: Origin and conservation. *Grassland Science in Europe*, *10*, 14.
- Pereira, H. M., & Navarro, L. M. (2015). *Rewilding European landscapes*. Springer International Publishing.
- Pöyry, J., Luoto, M., Paukkunen, J., Pykälä, J., Raatikainen, K., & Kuussaari, M. (2006). Different responses of plants and herbivore insects to a gradient of vegetation height: An indicator of the vertebrate grazing intensity and successional age. *Oikos*, *115*, 401–412.
- Queiroz, C., Beilin, R., Folke, C., & Lindborg, R. (2014). Farmland abandonment: Threat or opportunity for biodiversity conservation? A global review. *Frontiers in Ecology and the Environment*, *12*, 288–296.
- Rohatgi, A. (2021). *WebPlotDigitizer—Extract data from plots, images, and maps: Version 4.5*. <https://automeris.io/WebPlotDigitizer/>
- Rossetti, M. R., Tschamtkke, T., Aguilar, R., & Batáry, P. (2017). Responses of insect herbivores and herbivory to habitat fragmentation: A hierarchical meta-analysis. *Ecology Letters*, *20*, 264–272.
- Rudmann-Maurer, K., Weyand, A., Fischer, M., & Stöcklin, J. (2008). The role of landuse and natural determinants for grassland vegetation composition in the swiss Alps. *Basic and Applied Ecology*, *9*, 494–503.
- Sauermost, R., Freudig, D., Bonk, M., Sendtko, A., Genaust, H. & Gack, C. (Eds.). (1999). *Lexikon der Biologie*. Spektrum Akademischer Verlag, Heidelberg.
- Schils, R. L. M., Bufo, C., Rhymer, C. M., Francksen, R. M., Klaus, V. H., Abdalla, M., et al. (2022). Permanent grasslands in Europe: Land use change and intensification decrease their multifunctionality. *Agriculture, Ecosystems & Environment*, *330*, 107891.
- Schoof, N., Luick, R., Beaufoy, G., Jones, G., Einarsson, P., Ruiz, J., Stefanova, V., Fuchs, D., Windmaißer, T., Hötter, H., Jeromin, H., Nickel, H., Schumacher, J., & Ukhanova, M. (2019). *Grünlandschutz in Deutschland: Treiber der Biodiversität, Einfluss von Agrarumwelt- und Klimamaßnahmen, Ordnungsrecht, Molkerwirtschaft und Auswirkungen der Klima- und Energiepolitik*. BfN Skript. https://www.hs-rottenburg.net/fileadmin/user_upload/Forschung/Forschungsprojekte/Management/GAPGRUEN/BfN_Skript_539.pdf
- Shugart, H. H. (2013). Succession, phenomenon of. In *Encyclopedia of biodiversity*. Academic Press.
- Skórka, P., Settele, J., & Woyciechowski, M. (2007). Effects of management cessation on grassland butterflies in southern Poland. *Agriculture, Ecosystems & Environment*, *121*, 319–324.
- Slodowicz, D., Durbecq, A., Ladouceur, E., Eschen, R., Humbert, J., & Arlettaz, R. (2023). The relative effectiveness of different grassland restoration methods: A systematic literature search and meta-analysis. *Ecological Solutions and Evidence*, *4*, e12221.
- Spohn, M., Golte-Bechtle, M. & Spohn, R. (2015). *Was blüht denn da? Kosmos Naturführer*. 2nd edn. Kosmos.
- Steffan-Dewenter, I., & Leschke, K. (2003). Effects of habitat management on vegetation and above-ground nesting bees and wasps of orchard meadows in Central Europe. *Biodiversity and Conservation*, *12*, 1953–1968.
- Sterne, J. A., & Harbord, R. M. (2004). Funnel plots in meta-analysis. *The Stata Journal*, *4*, 127–141.
- Sterne, J. A. C., Sutton, A. J., Ioannidis, J. P. A., Terrin, N., Jones, D. R., Lau, J., Carpenter, J., Rücker, G., Harbord, R. M., Schmid, C. H., Tetzlaff, J., Deeks, J. J., Peters, J., Macaskill, P., Schwarzer, G., Duval, S., Altman, D. G., Moher, D., & Higgins, J. P. T. (2011). Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ*, *343*, 4002.
- Stoate, C., Báldi, A., Beja, P., Boatman, N. D., Herzog, I., van Doorn, A., & Báldi, A. (2009). Ecological impacts of early 21st century agricultural change in Europe—a review. *Journal of Environmental Management*, *91*, 22–46.
- Tälle, M., Deák, B., Poschod, P., Valkó, O., Westerberg, L., & Milberg, P. (2016). Grazing vs. mowing: A meta-analysis of biodiversity benefits for grassland management. *Agriculture, Ecosystems & Environment*, *222*, 200–212.
- Tanneberger, F., Bellebaum, J., Fartmann, T., Haferland, H.-J., Helmecke, A., Jehle, P., & Bellebaum, J. (2008). Rapid deterioration of aquatic warbler *Acrocephalus paludicola* habitats at the western margin of the breeding range. *Journal für Ornithologie*, *149*, 105–115.
- Tanneberger, F., Birr, F., Couwenberg, J., Kaiser, M., Luthardt, V., Neger, M., & Birr, F. (2022). Saving soil carbon, greenhouse gas emissions, biodiversity and the economy: Paludiculture as sustainable land use option in German fen peatlands. *Regional Environmental Change*, *22*, 69.
- Tanneberger, F., Flade, M., Preiksa, Z., & Schröder, B. (2010). Habitat selection of the globally threatened aquatic warbler *Acrocephalus paludicola* at the western margin of its breeding range and implications for management. *Ibis*, *152*, 347–358.
- Tonelli, M., Verdú, J. R., & Zunino, M. (2018). Effects of the progressive abandonment of grazing on dung beetle biodiversity: Body size matters. *Biodiversity and Conservation*, *27*, 189–204.
- Valkó, O., Török, P., Matus, G., & Tóthmérész, B. (2012). Is regular mowing the most appropriate and cost-effective management maintaining diversity and biomass of target forbs in mountain hay meadows? *Flora—Morphology, Distribution, Functional Ecology of Plants*, *207*, 303–309.
- Valkó, O., Venn, S., Žmihorski, M., Biurrun, I., Labadessa, R., & Loos, J. (2018). The challenge of abandonment for the sustainable management of Palaearctic natural and semi-natural grasslands. *Hacquetia*, *17*, 5–16.
- Verhulst, J., Báldi, A., & Kleijn, D. (2004). Relationship between land-use intensity and species richness and abundance of birds in Hungary. *Agriculture, Ecosystems & Environment*, *104*, 465–473.
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, *36*, 1–48.

- Vrahnakis, M., Janišová, M., Rüsina, S., Török, P., Venn, S., & Dengler, J. (2013). *The European dry grassland group (EDGG): Stewarding Europe's most diverse habitat type*. http://www.bayceer.uni-bayreuth.de/bayceer/de/pub/pub/120154/JD184_Vrahnakis_et_al_2013_Steppenrasenband.pdf
- Walcher, R., Hussain, R. I., Sachslehner, L., Böhner, A., Jernej, I., Zaller, J. G., Arnberger, A., & Frank, T. (2019). Long-term abandonment of mountain meadows affect bumblebees, true bugs and grasshoppers: A case study in the Austrian Alps. *Applied Ecology and Environmental Research*, 17, 5887–5908.
- WallisDeVries, M. F., Poschlod, P., & Willems, J. H. (2002). Challenges for the conservation of calcareous grasslands in northwestern Europe: Integrating the requirements of flora and fauna. *Biological Conservation*, 104, 265–273.
- Wiezik, M., Wieziková, A., & Svitok, M. (2011). Vegetation structure, ecological stability, and low-disturbance regime of abandoned dry grasslands support specific ant assemblages in Central Slovakia. *Tuexenia*, 31, 301–315.
- Wilson, J. B., Peet, R. K., Dengler, J., & Pärtel, M. (2012). Plant species richness: The world records. *Journal of Vegetation Science*, 23, 796–802.
- Woodward, F. I., Lomas, M. R., & Kelly, C. K. (2004). Global climate and the distribution of plant biomes. *Philosophical Transactions of The Royal Society B Biological Sciences*, 359, 1465–1476.
- Xiong, S., & Nilsson, C. (1999). The effects of plant litter on vegetation: A meta-analysis. *Journal of Ecology*, 87, 984–994.
- Zeiss, R., Eisenhauer, N., Orgiazzi, A., Rillig, M., Buscot, F., Jones, A., & Eisenhauer, N. (2022). Challenges of and opportunities for protecting European soil biodiversity. *Conservation Biology*, 36, e13930.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Elliott, T., Thompson, A., Klein, A.-M., Albert, C., Eisenhauer, N., Jansen, F., Schneider, A., Sommer, M., Straka, T., Settele, J., Sporbert, M., Tanneberger, F., & Mupepele, A.-C. (2023). Abandoning grassland management negatively influences plant but not bird or insect biodiversity in Europe. *Conservation Science and Practice*, 5(10), e13008. <https://doi.org/10.1111/csp2.13008>