



Reverse arthroplasty compared to hemiarthroplasty and open reduction and internal fixation for displaced proximal humerus fracture in patients above 60: a Bayesian network meta-analysis

Filippo Migliorini^{1,2,3} · Giuliano Sammaria⁴ · Luise Schäfer⁵ · Michael Memminger² · Francesco Simeone² · Nicola Maffulli^{6,7,8}

Received: 16 January 2025 / Accepted: 6 September 2025 / Published online: 18 October 2025

© The Author(s) 2025

Abstract

Introduction The present Bayesian network meta-analysis compared reverse total shoulder arthroplasty (rTSA) to hemiarthroplasty (HA) and open reduction and internal fixation (ORIF) for displaced proximal humeral fractures (PHF) in patients above 60.

Methods This study was conducted according to the PRISMA extension statement for reporting systematic reviews incorporating network meta-analyses of healthcare interventions. In December 2024, PubMed, Web of Science, and Embase databases were accessed. No time constraint was set for the search. All clinical studies were accessed comparing rTSA, ORIF and HA for displaced PHF in patients older than 60. Only studies which compared at least two of the interventions of interest were eligible when they reported a minimum of 12 months of follow-up. Two-, three-, and four-part displaced fractures and fractures with a head-splitting component were considered.

Results Data from 878 procedures were collected. 72% (631 of 878 patients) were women. The mean length of the follow-up was 33.8 ± 14.8 months. The mean age of the patients was 73.4 ± 4.6 years. Between groups, comparability was found in the mean age, the ratio of men to women, the length of the follow-up, and the time elapsed from injury to the procedure. The rTSA demonstrated the lowest rate of complications, followed by HA and ORIF. The rTSA demonstrated the lowest rate of revision, followed by HA and ORIF. Given the limited and heterogeneous data, only complications and revision rates were analysed in the network meta-analysis; functional outcomes were discussed narratively.

Conclusion In patients above 60 with displaced PHF, rTSA was associated with a lower complication and revision rate than ORIF and HA.

Level of evidence Level IV, systematic review and meta-analysis.

Keywords Displaced shoulder fracture · Arthroplasty · Hemiarthroplasty · Reverse total shoulder arthroplasty · rTSA · Open reduction and internal fixation · ORIF · Elderlies · Aged

Introduction

Proximal humeral fractures (PHF) account for up to 6% of all adult fractures, representing the third most common type of fracture, particularly in older women [1–7]. Additional risk factors are diabetes, epilepsy, depression, and dementia [8]. Despite PHF being common [9, 10], the most appropriate treatment remains controversial [11, 12], with some level I investigations recommending conservative management

of three- and four-part fractures [13, 14]. Age, bone quality, comorbidities, instability, general condition of the patient [15–20], and the type of fracture are the most important prognostic factors [21, 22]. In patients not suitable for surgery and minimally displaced or hairline fractures, a nonoperative approach is advocated [23–26]. Surgery is, however, still performed for displaced and comminuted fractures [21, 27]. Open reduction and internal fixation (ORIF), hemiarthroplasty (HA), and reverse total shoulder arthroplasty

(rTSA) are the most common surgical procedures for PHF in older adults [28–30].

However, in elderly patients with displaced PHF, the choice among ORIF, HA, and rTSA remains clinically debated. ORIF is typically preferred in less comminuted fractures where anatomical reduction is feasible, although outcomes may be compromised by osteoporotic bone and the risk of avascular necrosis [31–35]. HA was introduced to address these limitations, but it depends on tuberosity healing and rotator cuff integrity, which are often impaired in older patients [28, 36]. More recently, rTSA has emerged as a reliable option that bypasses these anatomical constraints, offering predictable functional outcomes even without tuberosity repair [2, 37–39]. However, questions remain regarding its precise indications and long-term results [40, 41]. These treatment considerations underscore the need for a comprehensive comparison across all three modalities in this patient population.

Previous meta-analyses have provided valuable insights into selected comparisons among rTSA, HA, and ORIF. However, a comprehensive synthesis encompassing all three modalities remains lacking [42, 43]. In addition, several studies have been published since the most recent review by Suroto et al. [43], reinforcing the need for an updated and methodologically advanced appraisal of the literature. Accordingly, a Bayesian network meta-analysis was conducted to compare ORIF, HA, and rTSA for treating displaced PHF in older adults, integrating direct and indirect evidence. By focusing on complication and revision rates, this study aims to support contemporary clinical decision-making in the surgical management of PHF. The authors hypothesised that rTSA would result in lower complication and revision rates than ORIF and HA.

Methods

Eligibility criteria

All the clinical studies published in peer-reviewed journals investigating rTSA, ORIF, and HA in the management of PHF were accessed. Only studies which compared at least two of the interventions of interest were eligible. Only studies that reported a minimum of 12 months of follow-up were considered. Studies that combined two or more interventions were not considered, nor were those which reported data on experimental pre- or post-operative procedures. Only studies which included patients with a minimum age of 60 years were eligible. According to the author's language capabilities, English, Italian, German, Spanish, and French articles were eligible. Only studies with levels I–IV of evidence, according to the Oxford Centre of Evidence-Based

Medicine [44], were considered. Reviews, opinions, letters, and editorials were not considered. Animals, in vitro, biomechanics, computational, and cadaveric studies were not eligible. Missing quantitative data on outcomes of interest warranted the exclusion of the study.

Search strategy

This study was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses: the 2020 PRISMA statement [45]. The PICOTD algorithm was preliminarily established:

- P (Problem): PHF in patients above 60;
- I (Intervention): surgical management;
- C (Comparison): rTSA, ORIF, HA;
- O (Outcomes): revisions and complications;
- T (Timing): minimum 12 months follow-up;
- D (Design): clinical study.

In December 2024, the following databases were accessed: PubMed, Web of Science, and Embase. No time constraint was set for the search. The Medical Subject Headings (MeSH) used for the database search are reported in Appendix A. No additional filters were used in the database search.

Selection and data collection

Two authors (**,**) independently performed the database search. All the resulting titles were screened by hand, and the abstract was accessed if suitable. The full text of the abstracts that matched the topic of interest was accessed. If the full text was not accessible or available, the article was not considered for inclusion. A cross-reference search of the bibliography of the full-text articles was also performed to identify additional relevant articles. Disagreements were debated and mutually solved by the authors. A third senior author (***) made the final decision in case of further disagreement.

Data items

Two authors (**,**) independently performed data extraction. The following data at baseline were extracted: author, year of publication, journal, length of follow-up, number of patients, mean age of patients with related conditions, and BMI. Data concerning complications and revisions were collected. Elective hardware removal was not considered a revision or complication. Data were extracted to Microsoft Office Excel version 16.72 (Microsoft Corporation, Redmond, USA).

Methodological quality assessment and quality of the recommendations

The risk of bias was evaluated following the guidelines in the Cochrane Handbook for Systematic Reviews of Interventions [46]. Two reviewers (**;**) independently assessed the risk of bias in the extracted studies. Disagreements were solved by a third senior author (**). Nonrandomised controlled trials (non-RCTs) were evaluated using the Risk of Bias in Nonrandomised Studies of Interventions (ROBINS-I) tool [47]. Seven domains of potential bias in non-RCTs were assessed. Possible confounding and the nature of patient selection before the start of the comparative intervention are assessed by two domains. A further domain assesses bias in the classification during the intervention. The final four domains assess the methodological quality after the intervention comparison has been implemented and relate to deviations from previously intended interventions, missing data, erroneous measurement of outcomes, and bias in the selection of reported outcomes. The figure of the ROBINS-I was elaborated using the Robvis Software (Risk-Of-Bias VISualization, Riskofbias.info, Bristol, UK) [48].

Synthesis methods

The main author (**) performed the statistical analyses following the recommendations of the Cochrane Handbook for Systematic Reviews of Interventions [49]. Baseline demographics were assessed through the IBM SPSS (International Business Machines Corporation, Armonk, USA) software. Mean and standard deviation were used for continuous variables, and frequency (events/ observations) for binary endpoints. Analysis of variance (ANOVA) was used to assess baseline comparability, with values of $P > 0.1$ considered satisfactory. The network meta-analyses were conducted using the STATA Software/MP, version 14.1 (StataCorporation, College Station, Texas, USA). The STATA routine for Bayesian hierarchical random-effects model analysis was used. The Log Odd Ratio (LOR) effect measure was adopted to analyse dichotomic data. The analyses of complications and revisions were conducted in a Bayesian hierarchical framework, which accommodates treatment arms with zero events without the need for continuity correction. The overall inconsistency was evaluated through the equation for global linearity via the Wald test. If $P_{\text{Wald}} > 0.5$, the null hypothesis could not be rejected, and the consistency assumption could be accepted at the overall level of each treatment. Both confidence (CI) and percentile (PrI) intervals were 95%. Edge and interval plots were obtained and evaluated. The funnel plot of each comparison was performed to assess data dispersion. The funnel plot of the most commonly reported outcome was used to

determine the risk of publication bias. Egger's linear regression was performed to assess asymmetry in the funnel plot, with values of $P_{\text{Egger}} < 0.05$ indicating statistically significant asymmetry.

Results

Study selection

The systematic literature search revealed a total of 801 studies. Of them, 498 were identified as duplicates and therefore excluded. Screening the abstracts of the remaining 303 investigations for eligibility resulted in excluding an additional 274 articles. The detailed reasons that led to exclusion were the following: study type and design ($N=113$), low level of evidence ($N=36$), not comparing at least two of the interventions of interest (rTSA and ORIF and HA) ($N=43$), combining two or more of the interventions ($N=22$), reporting data on experimental pre- or post-operative procedures ($N=17$), patients studied aged less than 60 years ($N=21$), follow-up shorter than 12 months ($N=16$), and language limitations ($N=6$). A further 17 studies missed quantitative data under the outcomes of interest and were not considered. In conclusion, 12 clinical trials were selected for inclusion in the present investigation. The study design was prospective in three studies, and nine had a retrospective study design. The results of the literature search are shown in Fig. 1.

Risk of publication bias

Egger's test of the endpoint complications ($P_{\text{Egger}}=0.5$) and revisions ($P_{\text{Egger}}=0.6$) found no statistically significant asymmetry, indicating a low risk of bias at the overall publication level (Fig. 2).

Methodological quality assessment

The ROBINS-I was applied to investigate the risk of bias for all articles included in the present study, as only non-RCTs were reviewed. No major concerns were present in most studies, and only a few had minor concerns, resulting in a predominantly low-risk rating in this domain. The lack of participant data for a defined follow-up period led to a serious risk of selection bias in one of the included studies. Other serious risks of bias were identified in one study because of deviations from the planned intervention and in another because of concerns about outcome measurement. Bias during the intervention was mostly low in the remaining studies, as were the domains assessed for risk of bias after the intervention, due to missing data and the reported

Fig. 1 PRISMA flow chart of the literature search





Fig. 2 Funnel plot endpoint complications (*left*) and revisions (*right*)

outcome selection. Given the broadly acceptable quality of the included studies, the overall risk of bias was low to moderate (Fig. 3).

Study characteristics and results of individual studies

Data from 878 procedures were collected. 72% (631 of 878 patients) were women. The mean length of the follow-up was 33.8±14.8 months. The mean age of the patients was 73.4±4.6 years. The generalities and demographics of the included studies are shown in Table 1. All pooled data from complications for the analysis of each procedure are reported alphabetically in Appendix B.

Baseline comparability

Comparability was found between groups regarding mean age, men:women ratio, length of follow-up, and time elapsed from injury to the index procedure (Table 2).

Synthesis of results

rTSA demonstrated the lowest rate of complications, followed by hemiarthroplasty and ORIF (Fig. 4). The equation for global linearity found no statistically significant inconsistency ($P_{Wald}=0.7$).

rTSA demonstrated the lowest rate of revision, followed by hemiarthroplasty and ORIF (Fig. 5). The equation for global linearity found no statistically significant inconsistency ($P_{Wald}=0.7$).

Discussion

According to the main findings of the present study, for patients above 60 with PHF in whom surgery was performed, rTSA was associated with a lower rate of complications and revision compared to ORIF and HA. These results contribute to the ongoing clinical discussion regarding the optimal surgical strategy in this population.

Osteosynthesis with preservation of the humeral head is recommended when adequate reduction can be obtained [31, 35]. However, avascular necrosis might occur, given the terminal vascularisation of the humeral head. Moreover, given the relatively high complication rate, alternative surgical options, such as arthroplasty and non-operative management, have been advocated [32–34]. ORIF performs better than hemiarthroplasty in less comminuted fractures, leading to better outcomes and range of motion with similar complication rates, improving the bending and torsion resistance [50, 59]. However, in comminuted PHF, HA was associated with lower pain on the VAS than ORIF [57, 60, 61]. Feng et al. [52] compared ORIF in adults (n=24) versus ORIF in




Fig. 3 The ROBINS-I of non-RCTs

Table 1 Generalities and patient baseline of the included studies

Author	Journal name	Design	Groups	Follow-up (month)	Patients (n)	Mean age	Women (n)
Cai et al., 2012 [50]	<i>Orthopedics</i>	Prospective	ORIF	24.3	13	71.1	11
			HA	24.3	19	72.4	16
Chalmers et al., 2014 [51]	<i>J Shoulder Elbow Surg</i>	Retrospective	ORIF	36.0	9	71.0	7
			HA	58.8	9	72.0	7
			rTSA	14.4	9	77.0	7
Feng et al., 2023 [52]	<i>Heliyon</i>	Retrospective	ORIF	37.4	36	71.3	29
			HA	56.1	10	73.5	7
Fraser et al., 2020 [37]	<i>J Bone Joint Surg Am</i>	Prospective	ORIF	24.0	60	74.7	52
			rTSA	24.0	64	75.7	59
Giardella et al., 2017 [38]	<i>Muscles Ligaments Tendons J</i>	Retrospective	ORIF	40.0	23	72.1	16
			rTSA	24.0	21	77.2	18
Klug et al., 2020 [39]	<i>J Shoulder Elbow Surg</i>	Retrospective	ORIF	49.0	30	72.5	5
			rTSA	38.0	30	73.9	5
Lanzetti et al., 2023 [53]	<i>Clin Orthop Relat Res</i>	Prospective	ORIF	53.0	66	73.0	29
			rTSA	53.0	72	76.0	40
Luciani et al., 2022 [54]	<i>Musculoskelet Surg</i>	Retrospective	ORIF	40.0	26	73.0	25
			rTSA	33.4	22	75.5	20
Muller et al., 2022 [55]	<i>Arch Orthop Trauma Surg</i>	Retrospective	ORIF	52.5	63	71.4	46
			rTSA	33.5	40	76.8	34
Ott et al., 2022 [56]	<i>Ota Int</i>	Retrospective	ORIF		90	81.9	76
			rTSA		71	82.3	56
Peker et al., 2022 [57]	<i>J Pak Med Assoc</i>	Retrospective	ORIF	18.3	30	60.0	14
			HA	18.7	18	67.3	11
Samborski et al., 2022 [58]	<i>J Shoulder Elbow Surg</i>	Retrospective	ORIF	12.0	23	67.1	18
			rTSA	12.0	24	77.3	23

ORIF open reduction and internal fixation; rTSA reverse total shoulder arthroplasty; HA hemiarthroplasty

Table 2 Baseline comparability (FU: follow-up; ORIF: open reduction and internal fixation; rTSA: reverse total arthroplasty)

Endpoint	Hemiarthroplasty (n=56)	ORIF (n=469)	rTSA (n=353)	P
Last FU (month)	39.5±20.9	35.1±13.9	29.0±13.4	0.3
Mean age	71.3±2.7	71.6±5.0	76.9±2.3	0.1
Women (%)	73% (41 of 56)	70% (328 of 469)	76% (262 of 353)	0.5
Time to procedure (days)	11.4±8.0	5.1±3.0	5.0±1.8	0.09

elderly patients (n=36) versus HA (n=10) in patients with locked fracture-dislocations of the proximal humerus. For patients younger than 60 years, ORIF might be optimal, whereas for patients older than 60 years, both ORIF and HA provided similar results [52]. However, ORIF was associated with a higher rate of complications [52].

rTSA might promote greater joint function than HA and ORIF [2, 38]. rTSA results in greater function than ORIF in bifocal extra-articular and intra-articular PHFs in




Fig. 4 Edge and interval plot of the comparison: complication




Fig. 5 Edge and interval plot of the comparison: revision

124 patients above 60 [37]. These findings are supported by previous clinical investigations, which recommended rTSA in older adults with dislocated PHFs, especially for its lower complication rate and revision [39, 54, 55, 58]. The superiority of rTSA to ORIF and HA was also confirmed in previous meta-analyses [37, 42, 43, 62]. However, there are no clear indications for rTSA [40, 41]. rTSA might be performed in non-reconstructable head-splitting fractures, fractures devoid of all soft tissue attachments, severe valgus-impacted fractures with disruption of the medial periosteal hinge, and displaced multi-fragmentary fractures with delayed presentation (more than four weeks from injury) [63–65]. HA was introduced to overcome the limitations of ORIF in patients with complex PHFs who required humeral head excision [28, 36]. However, HA is becoming increasingly less common, given the introduction of innovative locking plates and the good outcomes in rTSA [28, 66, 67]. HA is limited to patients with comminuted head fractures with intra-articular involvement and intact glenoid [28, 68, 69]. Instability, rotator cuff dysfunction, and tuberosity non-union leading to functional deficits represent possible complications of HA. In addition, the results of the present study indicate that HA is associated with a greater rate of complications and revision compared to rTSA. On the other hand, in patients with good shoulder function without advanced osteoarthritis and glenoid damage, HA spares the glenoid instrumentation, shortens surgical time, and reduces surgery-related costs.

Previous evidence is conflicting. Gallinet et al. [42] conducted a systematic review and meta-analysis of 22 studies comparing rTSA and HA in elderly patients with displaced proximal humeral fractures. While rTSA demonstrated superior functional outcomes in forward flexion and abduction, HA was associated with a lower complication rate [42]. The reoperation rate was comparable between groups,

whereas the revision rate (implant exchange) was higher after HA [42]. Suroto et al. [43] performed a meta-analysis of six comparative studies evaluating rTSA versus ORIF for three- and four-part fractures. Their results have shown that rTSA was associated with improved forward flexion and constant scores but also a higher complication rate compared to ORIF during early follow-up [43]. In contrast, the present analysis suggests that rTSA is associated with a lower rate of complications and revisions than ORIF and HA, particularly in older adults. In addition to these surgical outcomes, rTSA also tended to be associated with improved functional recovery, as reflected by higher Constant and ASES scores and lower VAS pain scores in several studies. A formal statistical comparison was not feasible given the inconsistent reporting and missing measures of dispersion. The discrepancy with previous meta-analyses may be attributable to differences in inclusion criteria, temporal shifts in surgical practice, and the incorporation of more recent evidence in the present study. Furthermore, applying a Bayesian network model instead of conventional pairwise comparisons may have contributed to the variation in findings by integrating direct and indirect evidence.

This study has several limitations, including the limited number of included studies, relative sample size, surgical techniques and type of implants, and the retrospective nature of most studies. The length of the follow-up was variable. Although several studies reported data on the ROM, the reporting was inconsistent across studies regarding completeness, measurement methods (e.g., active vs. passive ROM), and availability of statistical dispersion (e.g., standard deviations). In many cases, only partial ROM data were reported, but not all movements (flexion, abduction, external rotation) were consistently assessed across all surgical groups. Given these limitations, comparing the ROM was not possible in a network modality. Given

the variability in the definition of complications and their reporting across studies, a standardised categorisation with relative risk calculation and network comparison was not feasible without introducing bias. While some studies did report revision procedures in detail, others provided only general information, and definitions varied considerably. In addition, the complication rate was mainly low, which impaired the Bayesian model analysis in detecting a ranking of interventions. Therefore, a structured comparison of revision types was not performed. In some studies, the short follow-up might jeopardise the detection of long-term complications, especially regarding avascular necrosis or loosening of the prosthesis. Most studies did not specify the type of fractures or did not refer to the international classification; therefore, additional subgroup analyses according to the fracture classification were not possible. The present Bayesian network meta-analysis did not consider hardware removal. Hardware removal might be necessary for patients with stiffness, screw migration, or mechanical impingement, which represent surgical complications. On the other hand, hardware removal might follow patients' wishes and the healthcare system or health insurance indications, which is not a surgical complication. However, since most studies did not report information on the surgical indication, hardware removal was not considered for analysis. Despite this limitation, ORIF showed the highest complication rate, irrespective of hardware removal. Many included studies did not report clinical data or reported patient-reported outcome measures (PROMs) without any statistical measures of dispersion (e.g. standard deviation); therefore, given the lack of exhaustive data available for inclusion, a comparison of the clinical outcome between the three techniques could not be conducted in a network fashion. Future studies should overcome this limitation. The rTSA group has a shorter follow-up length than the other groups. Although these data were not statistically significant, they might induce an underestimation of the rate of complications of rTSA. In addition, one article did not report information on the length of the follow-up. Given the small number of included studies, the power to detect asymmetry of Egger's test and funnel plots is limited. Additional studies should evaluate patients according to internationally accepted classifications to refine indications and outcomes and provide longer follow-ups. Future studies should also assess which patients are suitable for conservative management or surgical treatment, integrating the advantage of current developments in artificial intelligence tools into patient care pathways [70].

Conclusion

In patients above 60 with PHF, rTSA was associated with a lower rate of complications and revision compared to ORIF and HA. Future investigations are required to validate these results on a larger scale and elaborate internationally accepted guidelines and therapeutic algorithms for managing PHF in this age group.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00402-025-06067-5>.

Author contributions Filippo Migliorini: conception and design, statistical analysis, drafting; Nicola Maffulli: supervision, revision; Giuliano Sammaria: writing, literature search, data extraction, risk of bias assessment; Luise Schäfer: literature search, data extraction, risk of bias assessment; Michael Memminger: supervision; Francesco Simeone: drafting. All authors have agreed to the final version to be published and agree to be accountable for all aspects of the work.

Funding Open Access funding enabled and organized by Projekt DEAL.

Data availability The datasets generated during and/or analysed during the current study are available throughout the manuscript.

Declarations

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bell JE, Leung BC, Spratt KF, Koval KJ, Weinstein JD, Goodman DC, Tosteson AN (2011) Trends and variation in incidence, surgical treatment, and repeat surgery of proximal humeral fractures in the elderly. *J Bone Joint Surg Am* 93(2):121–131. <https://doi.org/10.2106/JBJS.I.01505>
2. Court-Brown CM, Caesar B (2006) Epidemiology of adult fractures: a review. *Injury* 37(8):691–697. <https://doi.org/10.1016/j.injury.2006.04.130>
3. Passaretti D, Candela V, Sessa P, Gumina S (2017) Epidemiology of proximal humeral fractures: a detailed survey of 711 patients in a metropolitan area. *J Shoulder Elbow Surg* 26(12):2117–2124. <https://doi.org/10.1016/j.jse.2017.05.029>

4. Min W, Davidovitch RI, Tejwani NC (2012) Three-and four-part proximal humerus fractures: evolution to operative care. *Bull NYU Hosp Jt Dis* 70(1):25–34
5. Rose SH, Melton LJ 3rd, Morrey BF, Ilstrup DM, Riggs BL (1982) Epidemiologic features of humeral fractures. *Clin Orthop Relat Res* 168:24–30
6. Holloway KL, Bucki-Smith G, Morse AG, Brennan-Olsen SL, Kotowicz MA, Moloney DJ, Sanders KM, Korn S, Timney EN, Dobbins AG, Pasco JA (2015) Humeral fractures in South-Eastern Australia: epidemiology and risk factors. *Calcif Tissue Int* 97(5):453–465. <https://doi.org/10.1007/s00223-015-0039-9>
7. Lind T, Kroner K, Jensen J (1989) The epidemiology of fractures of the proximal humerus. *Arch Orthop Trauma Surg* 108(5):285–287. <https://doi.org/10.1007/BF00932316>
8. Roux A, Decroocq L, El Batti S, Bonneville N, Moineau G, Trojani C, Boileau P, de Peretti F (2012) Epidemiology of proximal humerus fractures managed in a trauma center. *Orthop Traumatol Surg Res* 98(6):715–719. <https://doi.org/10.1016/j.otsr.2012.05.013>
9. Han PF, Yang S, Wang YP, Hou XD, Li Y, Li XY (2022) Reverse shoulder arthroplasty vs. hemiarthroplasty for the treatment of osteoporotic proximal humeral fractures in elderly patients: a systematic review and meta-analysis update. *Exp Ther Med* 24(4):637. <https://doi.org/10.3892/etm.2022.11574>
10. Miquel J, Cassart E, Santana F, Martinez R, Valls L, Salomó-Domenech M, Torrens C (2024) Reverse shoulder arthroplasty or not for patients with displaced proximal humeral fractures: a randomized controlled trial. *J Shoulder Elbow Surg*. <https://doi.org/10.1016/j.jse.2024.02.023>
11. Neer CS 2nd (1970) Displaced proximal humeral fractures. II. Treatment of three-part and four-part displacement. *J Bone Joint Surg Am* 52(6):1090–1103
12. Neer CS 2nd (1970) Displaced proximal humeral fractures. I. Classification and evaluation. *J Bone Joint Surg Am* 52(6):1077–1089
13. Handoll HH, Elliott J, Thillemann TM, Aluko P, Brorson S (2022) Interventions for treating proximal humeral fractures in adults. *Cochrane Database Syst Rev* 6(6):CD000434. <https://doi.org/10.1002/14651858.CD000434.pub5>
14. Soler-Peiro M, Garcia-Martinez L, Aguilera L, Perez-Bermejo M (2020) Conservative treatment of 3-part and 4-part proximal humeral fractures: a systematic review. *J Orthop Surg Res* 15(1):347. <https://doi.org/10.1186/s13018-020-01880-7>
15. Olsson C, Petersson CJ (2006) Clinical importance of comorbidity in patients with a proximal humerus fracture. *Clin Orthop Relat Res* 442:93–99. <https://doi.org/10.1097/01.blo.0000194673.56764.0e>
16. Hertel R (2005) Fractures of the proximal humerus in osteoporotic bone. *Osteoporos Int* 16(Suppl 2):S65–72. <https://doi.org/10.1007/s00198-004-1714-2>
17. Masud S, Momtaz D, Betsch M, Migliorini F, Ghali A, Popa A, Gouveia K, Leroux T, Degen R, Khan M (2023) A comprehensive comparison and evaluation of surgical techniques for anterior shoulder instability: a Bayesian network meta-analysis. *J Shoulder Elbow Surg* 32(11):e531–e547. <https://doi.org/10.1016/j.jse.2023.07.004>
18. Poku D, Hassan R, Migliorini F, Maffulli N (2023) Efficacy of hydrodilatation in frozen shoulder: a systematic review and meta-analysis. *Br Med Bull* 147(1):121–147. <https://doi.org/10.1093/bmb/ldad018>
19. Silva ER, Maffulli N, Migliorini F, Santos GM, de Menezes FS, Okubo R (2022) Function, strength, and muscle activation of the shoulder complex in Crossfit practitioners with and without pain: a cross-sectional observational study. *J Orthop Surg Res* 17(1):24. <https://doi.org/10.1186/s13018-022-02915-x>
20. Neuhaus V, Swellingrebel CH, Bossen JK, Ring D (2013) What are the factors influencing outcome among patients admitted to a hospital with a proximal humeral fracture? *Clin Orthop Relat Res* 471(5):1698–1706. <https://doi.org/10.1007/s11999-013-2876-z>
21. Troiano E, Peri G, Calo I, Colasanti GB, Mondanelli N, Giannotti S (2023) A novel “7 sutures and 8 knots” surgical technique in reverse shoulder arthroplasty for proximal humeral fractures: tuberosity healing improves short-term clinical results. *J Orthop Traumatol* 24(1):18. <https://doi.org/10.1186/s10195-023-00697-4>
22. Yang J, Yang Z, Liu D, Lu Z, Tao C, Liu T (2023) Is an antero-medial minimally invasive approach for middle and distal third humeral fractures feasible? A cadaveric study and clinical case series. *J Orthop Traumatol* 24(1):7. <https://doi.org/10.1186/s10195-023-00684-9>
23. Patel AH, Wilder JH, Ofa SA, Lee OC, Savoie FH 3rd, O'Brien MJ, Sherman WF (2022) Trending a decade of proximal humerus fracture management in older adults. *JSES Int* 6(1):137–143. <https://doi.org/10.1016/j.jseint.2021.08.006>
24. Schumaier A, Grawe B (2018) Proximal humerus fractures: evaluation and management in the elderly patient. *Geriatr Orthop Surg Rehabil* 9:2151458517750516. <https://doi.org/10.1177/2151458517750516>
25. Kimmeyer M, Schmalz J, Rentschler V, Schieffer C, Macken A, Gerhardt C, Lehmann LJ (2023) Correct positioning of the calcar screw leads to superior results in proximal humerus fractures treated with carbon-fibre-reinforced polyetheretherketone plate osteosynthesis with polyaxial locking screws. *J Orthop Traumatol* 24(1):54. <https://doi.org/10.1186/s10195-023-00733-3>
26. Wu K, Lin T, Lee CH (2023) Intramedullary nailing versus cemented plate for treating metastatic pathological fracture of the proximal humerus: a comparison study and literature review. *J Orthop Traumatol* 24(1):45. <https://doi.org/10.1186/s10195-023-00721-7>
27. Fitschen-Oestern S, Behrendt P, Martens E, Finn J, Schiegnitz J, Borzиковsky C, Seekamp A, Weuster M, Lippross S (2020) Reversed shoulder arthroplasty for the treatment of proximal humerus fracture in the elderly. *J Orthop* 17:180–186. <https://doi.org/10.1016/j.jor.2019.08.029>
28. Baker HP, Gutbrod J, Strelzow JA, Maassen NH, Shi L (2022) Management of proximal humerus fractures in adults—a scoping review. *J Clin Med*. <https://doi.org/10.3390/jcm11206140>
29. Brorson S, Elliott J, Thillemann T, Aluko P, Handoll H (2022) Interventions for proximal humeral fractures: Key messages from a cochrane review. *Acta Orthop* 93:610–612. <https://doi.org/10.2340/17453674.2022.3495>
30. Iacobellis C, Berizzi A, Biz C, Camporese A (2015) Treatment of proximal humeral fractures with reverse shoulder arthroplasty in elderly patients. *Musculoskelet Surg* 99(1):39–44. <https://doi.org/10.1007/s12306-014-0331-2>
31. Bastian JD, Hertel R (2009) Osteosynthesis and hemiarthroplasty of fractures of the proximal humerus: outcomes in a consecutive case series. *J Shoulder Elbow Surg* 18(2):216–219. <https://doi.org/10.1016/j.jse.2008.09.015>
32. Olerud P, Ahrengart L, Ponzer S, Saving J, Tidermark J (2011) Internal fixation versus nonoperative treatment of displaced 3-part proximal humeral fractures in elderly patients: a randomized controlled trial. *J Shoulder Elbow Surg* 20(5):747–755. <https://doi.org/10.1016/j.jse.2010.12.018>
33. Rangan A, Gwilym S, Keding A, Corbacho B, Kottam L, Arundel C, Coleman E, DiMascio L, Hewitt C, Jones V, Kassam J, McDaid C, Mitchell N, Mott A, O'Carroll G, Tharmanathan P, Torgerson D (2023) Reverse shoulder arthroplasty versus hemiarthroplasty versus non-surgical treatment for older adults with acute 3- or 4-part fractures of the proximal humerus: study protocol for a randomised controlled trial (PROFHER-2: PROximal fracture of humerus evaluation by randomisation - trial number 2). *Trials* 24(1):270. <https://doi.org/10.1186/s13063-023-07259-3>

34. Sidor ML, Zuckerman JD, Lyon T, Koval K, Schoenberg N (1994) Classification of proximal humerus fractures: the contribution of the scapular lateral and axillary radiographs. *J Shoulder Elbow Surg* 3(1):24–27. [https://doi.org/10.1016/S1058-2746\(09\)80004-9](https://doi.org/10.1016/S1058-2746(09)80004-9)
35. Solberg BD, Moon CN, Franco DP, Paiement GD (2009) Locked plating of 3- and 4-part proximal humerus fractures in older patients: the effect of initial fracture pattern on outcome. *J Orthop Trauma* 23(2):113–119. <https://doi.org/10.1097/BOT.0b013e31819344bf>
36. Neer CS 2nd (2011) The classic: articular replacement for the humeral head. 1955. *Clin Orthop Relat Res* 469(9):2409–2421. <https://doi.org/10.1007/s11999-011-1944-5>
37. Fraser AN, Bjordal J, Wagle TM, Karlberg AC, Lien OA, Eilertsen L, Mader K, Apold H, Larsen LB, Madsen JE, Fjalestad T (2020) Reverse shoulder arthroplasty is superior to plate fixation at 2 years for displaced proximal humeral fractures in the elderly: a multicenter randomized controlled trial. *J Bone Joint Surg Am* 102(6):477–485. <https://doi.org/10.2106/JBJS.19.01071>
38. Giardella A, Ascione F, Mocchi M, Berlusconi M, Romano AM, Oliva F, Maradei L (2017) Reverse total shoulder versus angular stable plate treatment for proximal humeral fractures in over 65 years old patients. *Muscles Ligaments Tendons J* 7(2):271–278. <https://doi.org/10.11138/mltj/2017.7.2.271>
39. Klug A, Harth J, Hoffmann R, Gramlich Y (2020) Surgical treatment of complex proximal humeral fractures in elderly patients: a matched-pair analysis of angular-stable plating vs. reverse shoulder arthroplasty. *J Shoulder Elbow Surg* 29(9):1796–1803. <https://doi.org/10.1016/j.jse.2020.01.073>
40. Gomberawalla MM, Miller BS, Coale RM, Bedi A, Gagnier JJ (2013) Meta-analysis of joint preservation versus arthroplasty for the treatment of displaced 3- and 4-part fractures of the proximal humerus. *Injury* 44(11):1532–1539. <https://doi.org/10.1016/j.injury.2013.03.030>
41. Sobel AD, Shah KN, Paxton ES (2017) Fixation of a proximal humerus fracture with an intramedullary nail. *J Orthop Trauma* 31(Suppl 3):S47–S49. <https://doi.org/10.1097/BOT.0000000000000909>
42. Gallinet D, Ohl X, Decroocq L, Dib C, Valenti P, Boileau P, French Society for Orthopaedic S (2018) Is reverse total shoulder arthroplasty more effective than hemiarthroplasty for treating displaced proximal humerus fractures in older adults? A systematic review and meta-analysis. *Orthop Traumatol Surg Res* 104(6):759–766. <https://doi.org/10.1016/j.otsr.2018.04.025>
43. Suroto H, De Vega B, Deapsari F, Prajasari T, Wibowo PA, Samijo SK (2021) Reverse total shoulder arthroplasty (RTSA) versus open reduction and internal fixation (ORIF) for displaced three-part or four-part proximal humeral fractures: a systematic review and meta-analysis. *EFORT Open Rev* 6(10):941–955. <https://doi.org/10.1302/2058-5241.6.210049>
44. Howick J CI, Glasziou P, Greenhalgh T, Carl Heneghan, Liberati A, Moschetti I, Phillips B, Thornton H, Goddard O, Hodgkinson M (2011) The 2011 Oxford CEBM levels of evidence. Oxford Centre for evidence-based medicine available at <https://www.cebm.net/index.aspx?o=5653>
45. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, Chou R, Glanville J, Grimshaw JM, Hrobjartsson A, Lalu MM, Li T, Loder EW, Mayo-Wilson E, McDonald S, McGuinness LA, Stewart LA, Thomas J, Tricco AC, Welch VA, Whiting P, Moher D (2021) The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 372:n71. <https://doi.org/10.1136/bmj.n71>
46. Cumpston M, Li T, Page MJ, Chandler J, Welch VA, Higgins JP, Thomas J (2019) Updated guidance for trusted systematic reviews: a new edition of the cochrane handbook for systematic reviews of interventions. *Cochrane Database Syst Rev* 10:CD000142. <https://doi.org/10.1002/14651858.ED000142>
47. Sterne JA, Hernan MA, Reeves BC, Savovic J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT, Boutron I, Carpenter JR, Chan AW, Churchill R, Deeks JJ, Hrobjartsson A, Kirkham J, Juni P, Loke YK, Pigott TD, Ramsay CR, Regidor D, Rothstein HR, Sandhu L, Santaguida PL, Schunemann HJ, Shea B, Shrier I, Tugwell P, Turner L, Valentine JC, Waddington H, Waters E, Wells GA, Whiting PF, Higgins JP (2016) ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 355:i4919. <https://doi.org/10.1136/bmj.i4919>
48. McGuinness LA, Higgins JPT (2021) Risk-of-bias VISualization (robvis): an R package and Shiny web app for visualizing risk-of-bias assessments. *Res Synth Methods* 12(1):55–61. <https://doi.org/10.1002/jrsm.1411>
49. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA. *Cochrane Handbook for Systematic Reviews of Interventions* version 6.2. Cochrane 2021. Available from <https://www.cochrane.org/resources/levels-of-evidence/ocebml-levels-of-evidence> Accessed on Feb 2022.
50. Cai M, Tao K, Yang C, Li S (2012) Internal fixation versus shoulder hemiarthroplasty for displaced 4-part proximal humeral fractures in elderly patients. *Orthopedics* 35(9):e1340–1346. <https://doi.org/10.3928/01477447-20120822-19>
51. Chalmers PN, Slikker W 3rd, Mall NA, Gupta AK, Rahman Z, Enriquez D, Nicholson GP (2014) Reverse total shoulder arthroplasty for acute proximal humeral fracture: comparison to open reduction-internal fixation and hemiarthroplasty. *J Shoulder Elbow Surg* 23(2):197–204. <https://doi.org/10.1016/j.jse.2013.07.044>
52. Feng D, Jiang Y, Kang X, Song Z, Zhu Y, Zhang J, Zhang K, Wang Z (2023) Assessment of surgical management for locked fracture-dislocations of the proximal humerus in patients of different ages. *Heliyon* 9(6):e16508. <https://doi.org/10.1016/j.heliyon.2023.e16508>
53. Lanzetti RM, Gaj E, Berlinberg EJ, Patel HH, Spoliti M (2023) Reverse total shoulder arthroplasty demonstrates better outcomes than angular stable plate in the treatment of three-part and four-part proximal humerus fractures in patients older than 70 years. *Clin Orthop Relat Res* 481(4):735–747. <https://doi.org/10.1097/CORR.0000000000002480>
54. Luciani P, Procaccini R, Rotini M, Pettinari F, Gigante A (2022) Angular stable plate versus reverse shoulder arthroplasty for proximal humeral fractures in elderly patient. *Musculoskeletal Surg* 106(1):43–48. <https://doi.org/10.1007/s12306-020-00669-5>
55. Muller M, Greve F, Cronlein M, Zyskowski M, Pesch S, Biberthaler P, Kirchoff C, Beirer M (2022) Reconstruction or replacement? A challenging question in surgical treatment of complex humeral head fractures in the elderly. *Arch Orthop Trauma Surg* 142(11):3247–3254. <https://doi.org/10.1007/s00402-021-04124-3>
56. Ott N, Muller C, Jacobs A, Paul C, Wegmann K, Muller LP, Kabir K (2022) Outcome of geriatric proximal humeral fractures: a comparison between reverse shoulder arthroplasty versus open reduction and internal fixation. *OTA Int* 5(2 Suppl):e188. <https://doi.org/10.1097/OI9.0000000000000188>
57. Peker B, Polat AE, Carkci E, Senel A, Soydan C, Tuzuner T (2022) Functional outcomes and complication analysis of plate osteosynthesis versus hemiarthroplasty in three-part and four-part proximal humerus fractures. *J Pak Med Assoc* 72(1):57–61. <https://doi.org/10.47391/JPMA.1147>
58. Samborski SA, Haws BE, Karnyski S, Soles G, Gorczyca JT, Nicandri G, Voloshin I, Ketz JP (2022) Outcomes for type C proximal humerus fractures in the adult population: comparison of nonoperative treatment, locked plate fixation, and reverse

- shoulder arthroplasty. *JSES Int* 6(5):755–762. <https://doi.org/10.1016/j.jseint.2022.05.006>
59. Edwards SL, Wilson NA, Zhang LQ, Flores S, Merk BR (2006) Two-part surgical neck fractures of the proximal part of the humerus. A biomechanical evaluation of two fixation techniques. *J Bone Joint Surg Am* 88(10):2258–2264. <https://doi.org/10.2106/JBJS.E.00757>
 60. Amundsen A, Rasmussen JV, Olsen BS, Brorson S (2019) Low revision rate despite poor functional outcome after stemmed hemiarthroplasty for acute proximal humeral fractures: 2,750 cases reported to the Danish Shoulder Arthroplasty Registry. *Acta Orthop* 90(3):196–201. <https://doi.org/10.1080/17453674.2019.1597491>
 61. Valenti P, Aliani D, Maroun C, Werthel JD, Elkolti K (2017) Shoulder hemiarthroplasty for proximal humeral fractures: analysis of clinical and radiographic outcomes at midterm follow-up: a series of 51 patients. *Eur J Orthop Surg Traumatol* 27(3):309–315. <https://doi.org/10.1007/s00590-017-1927-7>
 62. Szerlip BW, Morris BJ, Edwards TB (2016) Reverse shoulder arthroplasty for trauma: when, where, and how. *Instr Course Lect* 65:171–179
 63. Robinson CM, Khan LA, Akhtar MA (2006) Treatment of anterior fracture-dislocations of the proximal humerus by open reduction and internal fixation. *J Bone Joint Surg Br* 88(4):502–508. <https://doi.org/10.1302/0301-620X.88B4.17195>
 64. Bastian JD, Hertel R (2008) Initial post-fracture humeral head ischemia does not predict development of necrosis. *J Shoulder Elbow Surg* 17(1):2–8. <https://doi.org/10.1016/j.jse.2007.03.026>
 65. Hertel R, Hempfing A, Stiehler M, Leunig M (2004) Predictors of humeral head ischemia after intracapsular fracture of the proximal humerus. *J Shoulder Elbow Surg* 13(4):427–433. <https://doi.org/10.1016/j.jse.2004.01.034>
 66. Kontakis G, Tosounidis T, Galanakis I, Megas P (2008) Prosthetic replacement for proximal humeral fractures. *Injury* 39(12):1345–1358. <https://doi.org/10.1016/j.injury.2008.09.003>
 67. Schultz BJ, Lowe DT, Egol KA, Zuckerman JD (2021) Shoulder hemiarthroplasty for proximal humerus fracture. *J Orthop Trauma* 35(Suppl 2):S3–S4. <https://doi.org/10.1097/BOT.0000000000002158>
 68. Shukla DR, McAnany S, Kim J, Overley S, Parsons BO (2016) Hemiarthroplasty versus reverse shoulder arthroplasty for treatment of proximal humeral fractures: a meta-analysis. *J Shoulder Elbow Surg* 25(2):330–340. <https://doi.org/10.1016/j.jse.2015.08.030>
 69. Wilfred AM, Akhter S, Horner NS, Aljedani A, Khan M, Alolabi B (2022) Outcomes and complications of distal humeral hemiarthroplasty for distal humeral fractures - a systematic review. *Shoulder Elbow* 14(1):65–74. <https://doi.org/10.1177/17585732211023100>
 70. Zeng GJ, Sheng X, Lie DTT (2023) Short- to medium-term outcomes and future direction of reverse shoulder arthroplasty: current concepts. *J ISAKOS* 8(6):398–403. <https://doi.org/10.1016/j.jisako.2023.10.005>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Filippo Migliorini^{1,2,3} · Giuliano Sammaria⁴ · Luise Schäfer⁵ · Michael Memminger² · Francesco Simeone² · Nicola Maffulli^{6,7,8}

✉ Filippo Migliorini
Filippo.Migliorini@uk-halle.de

Giuliano Sammaria
g.sammaria@unisa.it

Luise Schäfer
luiseschaefer83@gmail.com

Michael Memminger
michael.memminger@sabes.it

Francesco Simeone
francesco.simeone@sabes.it

Nicola Maffulli
nicola.maffulli@uniroma1.it

¹ Department of Trauma and Reconstructive Surgery, University Hospital in Halle, Martin-Luther University Halle-Wittenberg, 06097 Halle (Saale), Germany

² Department of Orthopaedic and Trauma Surgery, Academic Hospital of Bolzano (SABES-ASDAA), via Lorenz Böhler 5, 39100 Bolzano, Italy

³ Department of Life Sciences, Health, and Health Professions, Link Campus University, Via del Casale di San Pio V, 00165 Rome, Italy

⁴ University of Salerno, Fisciano, Italy

⁵ Department of Orthopaedic and Trauma Surgery, Eifelklinik St. Brigida, Kammerbruschstr. 8, 52152 Simmerath, Germany

⁶ Faculty of Medicine and Psychology, University “La Sapienza” of Rome, Rome, Italy

⁷ School of Pharmacy and Bioengineering, Keele University Faculty of Medicine, Stoke on Trent, ST4 7QB London, United Kingdom

⁸ Centre for Sports and Exercise Medicine, Queen Mary University of London, Barts and the London School of Medicine and Dentistry, Mile End Hospital, 275 Bancroft Road, E1 4DG London, United Kingdom