

Lichenes selecti exsiccati ex Herbario Universitatis Halensis – nos. 11–20

Regine STORDEUR

Abstract: STORDEUR, R. 2007: Lichenes selecti exsiccati ex Herbario Universitatis Halensis – nos. 11–20. *Schlechtendalia* **16**: 93–97.

The second issue of 'Lichenes selecti exsiccati ex Herbario Universitatis Halensis' containing nos. 11–20 is presented. All collections are from copper schist dumps of the 'Mansfelder Mulde' area (Saxony-Anhalt).

Zusammenfassung: STORDEUR, R. 2007: Lichenes selecti exsiccati ex Herbario Universitatis Halensis – Nr. 11–20. *Schlechtendalia* **16**: 93–97.

Mit den vorliegenden Nummern 11–20 wird die 2. Ausgabe von 'Lichenes selecti exsiccati ex Herbario Universitatis Halensis' vorgelegt. Die Belege stammen alle von Kupferschieferhalde der Mansfelder Mulde (Sachsen-Anhalt).

Key words: HAL, exsiccata, lichens, copper schist.

The second series of 'Lichenes selecti exsiccati' exclusively comprises lichens collected from copper schist, including various characteristic heavy metal lichens as well as some species with a wider ecological amplitude which may also occur on other rocks.

The beginning of the copper schist mining in the area around the town Mansfeld ('Mansfelder Revier') dates back to about the year 1200. The dumps used as sources for lichen exsiccata are, however, much younger, i.e., the oblong 'Zirkelschachthalde' near the town Klostermansfeld, raised up between 1895 and 1927, and the large trapezoidal dump of the 'Eduard shaft' south of the town Hettstedt, used between 1864 and 1910. Beside some gypsum, the two locations are characterised by being composed of copper schist, partly with considerable amounts of metals (copper, ferrum, manganese, molybdenum, nickel, lead, titanium, zinc and other metals usually occurring as sulphides), as well as Permian lime-stone, a relatively solid stone containing lime.

For the present issue of this exsiccata only lichens directly growing on copper schist have been collected, although a certain influence of lime cannot be excluded due to the edulcoration of particular components of decaying Permian lime-stone.

The area 'Mansfelder Mulde' is located at the rain protected side of the Harz Mountains characterised by a low precipitation between 480 and 500 mm. The average temperature per year is 8.6 °C, with maximums of 15 to 17 °C in July and minimums of 0 to –2 °C in January.

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

11. *Acarospora peliscypha* Th. Fr.

Nova Acta Regiae Soc. Sci. Upsal., Ser. 3., 3: 189 (1861).

GERMANY, Saxony-Anhalt, NNW Lutherstadt-Eisleben, S Thondorf, copper schist damp of the 'Zirkelschacht', ordnance map 4435/114, 51°34'36"–52°N/11°31'40"–49"E, alt. c. 220–255 m, on copper schist.

12 Nov. 2003

leg. M. Beckmann & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

12. *Acarospora smaragdula* (Wahlenb.) A. Massal.

Ric. auton. lich. crost.: 29 (1852).

GERMANY, Saxony-Anhalt, NNW Lutherstadt-Eisleben, S Thondorf, copper schist damp of the 'Zirkelschacht', ordnance map 4435/114, 51°34'36"–52°N/11°31'40"–49"E, alt. c. 220–255 m, on copper schist.

12 Nov. 2003

leg. M. Beckmann & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

13. *Buellia aethalea* (Ach.) Th. Fr.

Lichenogr. scand. 2: 604 (1874).

GERMANY, Saxony-Anhalt, NNW Lutherstadt-Eisleben, S Thondorf, copper schist damp of the 'Zirkelschacht', ordnance map 4435/114, 51°34'36"–52°N/11°31'40"–49"E, alt. c. 220–255 m, on copper schist.

5 June 2007

leg. E. Eisenhuth & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

14. Candelariella vitellina (Hoffm.) Müll. Arg.

Bull. Herb. Boissier 2: 11, 47 (1894).

GERMANY, Saxony-Anhalt, large copper schist dump of the ‘Eduard’ shaft Hettstedt, ordnance map 4435/114, 51°37'17"–31"N/11°31'02"–28"E, alt. 200–222 m, on copper schist.

15 Nov. 2006

leg. M. Schönbrodt & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

15. Lecanora muralis (Schreb.) Rabenh.

Deutschl. Krypt.-Fl. 2: 42 (1845).

GERMANY, Saxony-Anhalt, large copper schist dump of the ‘Eduard’ shaft Hettstedt, ordnance map 4435/114, 51°37'17"–31"N/11°31'02"–28"E, alt. 200–222 m, on copper schist.

15 Nov. 2006

leg. M. Schönbrodt & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

16. Lecanora stenotropa Nyl.

Flora 55: 251 (1872).

GERMANY, Saxony-Anhalt, NNW Lutherstadt-Eisleben, S Thondorf, copper schist dump of the ‘Zirkelschacht’, ordnance map 4435/114, 51°34'36"–52"N/11°31'40"–49"E, alt. c. 220–255 m, on copper schist.

5 June 2007

leg. E. Eisenhuth & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

17. Lecidea fuscoatra (L.) Ach.

Methodus, Sectio prior: 44 (1803).

GERMANY, Saxony-Anhalt, NNW Lutherstadt-Eisleben, S Thondorf, copper schist dump of the 'Zirkelschacht', ordnance map 4435/114, 51°34'36"–52°N/11°31'40"–49"E, alt. c. 220–255 m, on copper schist.

5 June 2007

leg. E. Eisenhuth & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

18. Lecidea inops Th. Fr.

Lichenogr. scand. 2: 501 (1874).

GERMANY, Saxony-Anhalt, large copper schist dump of the 'Eduard' shaft Hettstedt, ordnance map 4435/114, 51°37'17"–31"N/11°31'02"–28"E, alt. 200–222 m, on copper schist.

15 Nov. 2006

leg. M. Schönbrodt & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati
(ex Herbario Universitatis Halensis)

19. Scoliciosporum umbrinum (Ach.) Arnold

Flora 54: 50 (1871).

GERMANY, Saxony-Anhalt, NNW Lutherstadt-Eisleben, S Thondorf, copper schist dump of the 'Zirkelschacht', ordnance map 4435/114, 51°34'36"–52°N/11°31'40"–49"E, alt. c. 220–255 m, on copper schist.

5 June 2007

leg. E. Eisenhuth & R. Stordeur
det. R. Stordeur

R. Stordeur: Lichenes selecti exsiccati (ex Herbario Universitatis Halensis)

20. *Stereocaulon nanodes* Tuck.

Amer. J. Arts Sci., Ser. 2, 18: 201 (1859).

GERMANY, Saxony-Anhalt, large copper schist dump of the ‘Eduard’ shaft Hettstedt, ordnance map 4435/114, 51°37'17"–31"N/11°31'02"–28"E, alt. 200–222 m, on copper schist.

15 Nov. 2006

leg. M. Schönbrodt & R. Stordeur
det. R. Stordeur

Acknowledgements

I am grateful to M. Beckmann, E. Eisenhuth and M. Schönbrodt for their help to collect some specimens for the second issue (serie) of this exsiccata. I am much obliged to U. Braun for correcting the English text.

Address of the author

Regine Stordeur, Martin-Luther-Universität, Institut für Biologie, Geobotanik und Botanischer Garten, Neuwerk 21, D-06099 Halle (Saale), Germany.
(E-mail: regine.stordeur@botanik.uni-halle.de)